

2018 • 2019

Faculteit Industriële ingenieurswetenschappen
master in de industriële wetenschappen: biochemie

Masterthesis

Uranium Toxicity to the Freshwater Plant *Lemna minor* in an Environmentally Relevant Metal Mixture: Selection and Validation of Potential Uranium-Biomarker Genes

PROMOTOR :

dr. ir. Kristel SNIEGOWSKI

PROMOTOR :

dr. Eline SAENEN

COPROMOTOR :

prof. dr. Nele HOREMANS

Isabelle Van Dyck

Scriptie ingediend tot het behalen van de graad van master in de industriële wetenschappen: biochemie

Gezamenlijke opleiding UHasselt en KU Leuven



KU LEUVEN



KU LEUVEN

2018•2019

Faculteit Industriële ingenieurswetenschappen
master in de industriële wetenschappen: biochemie

Masterthesis

Uranium Toxicity to the Freshwater Plant *Lemna minor* in an Environmentally Relevant Metal Mixture: Selection and Validation of Potential Uranium-Biomarker Genes

PROMOTOR :

dr. ir. Kristel SNIÉGOWSKI

PROMOTOR :

dr. Eline SAENEN

COPROMOTOR :

prof. dr. Nele HOREMANS

Isabelle Van Dyck

Scriptie ingediend tot het behalen van de graad van master in de industriële wetenschappen: biochemie



KU LEUVEN

ACKNOWLEDGEMENTS

After a bridging program and a master year, the final step to graduation as an industrial engineer in Biochemistry, is the internship and master thesis. These two years have flown by and would not have been possible with the help of certain people. That is why I want to thank these people for their help and contributions.

First of all I want to thank my promotor dr. Eline Saenen to help and guide me during my thesis. Thank you for answering all of my questions and thoroughly read my thesis many times. I also want to thank prof. dr. Nele Horemans for the opportunity to perform this project within the Biosphere Impact Studies (BIS) group at SCK•CEN. It was a very interesting and educational subject at which I learned a lot. Additionally I would like to thank May Van Hees, Axel Van Gompel, Robin Nauts and Jean Wannijn to help me with the experimental work and for answering all the questions that came with it. Also thank you Axel, Robin and Jean for the great times and talks at the office. Furthermore I would like to thank all the other colleagues at BIS and the fellow trainees for the nice talks and the good atmosphere. Also a sincere thank you to my university promotor dr. ir. Kristel Sniegowski for the opportunity to perform my internship at SCK•CEN, to read my thesis and to give feedback during the process.

I also would like to thank my parents who gave me the opportunity to start this education after my bachelor's degree. They believed in me during these past two years and I am grateful for that. And last but not least I would like to thank my classmates that also became very close friends. Thank you Anneleen, Demi, Julie and Wouter for the amazing two years and the years still to come!

TABLE OF CONTENTS

| | |
|---|-----------|
| ACKNOWLEDGEMENTS | 1 |
| LIST OF TABLES | 5 |
| LIST OF FIGURES | 7 |
| ABBREVIATIONS | 9 |
| ABSTRACT | 11 |
| ABSTRACT IN DUTCH | 13 |
| 1 INTRODUCTION | 15 |
| 2 ENVIRONMENTAL POLLUTION | 17 |
| 2.1 Uranium | 17 |
| 2.2 Environmentally relevant metal mixture | 18 |
| 2.3 <i>Lemna minor</i> | 20 |
| 2.4 Biomarkers | 21 |
| 3 METHODS AND MATERIALS | 23 |
| 3.1 <i>L. minor</i> growth and exposure to uranium and metal mixture | 23 |
| 3.2 Specific growth rate and growth inhibition | 24 |
| 3.3 Primer development and optimization | 25 |
| 3.4 Measurements on molecular level | 27 |
| 3.4.1 Gene expression | 27 |
| 3.4.2 Comet assay | 28 |
| 3.5 Statistical analysis | 30 |
| 4 RESULTS | 31 |
| 4.1 Primer selection and optimization | 31 |
| 4.2 Primer selectivity | 31 |
| 4.3 7 days exposure experiment | 35 |
| 4.4 Time kinetics experiment | 38 |
| 4.5 Summary of the gene expression results | 44 |
| 4.6 Comet assay | 45 |
| 5 DISCUSSION | 47 |
| 5.1 Growth effects induced by exposure to U and the metal mixture | 47 |
| 5.2 Gene expression of potential U-biomarkers | 48 |
| 5.3 Comet assay optimization on <i>L. minor</i> | 49 |
| CONCLUSION | 51 |
| REFERENCES | 53 |
| APPENDIX A: COMPOSITION OF THE CONTAMINATION SOLUTION OF THE 7 DAYS EXPOSURE EXPERIMENT AND THE TIME KINETICS EXPERIMENT | 63 |
| APPENDIX B: SPECIFICITY TESTING OF POTENTIAL HOUSEKEEPING GENES FOR <i>L. MINOR</i> | 65 |
| APPENDIX C: SPECIFICITY TESTING OF POTENTIAL U-BIOMARKER GENES FOR <i>L. MINOR</i> | 71 |
| APPENDIX D: GROWTH OF <i>L. MINOR</i> PLANTS IN 7 DAYS EXPOSURE EXPERIMENT | 75 |
| APPENDIX E: GROWTH OF <i>L. MINOR</i> PLANTS IN TIME KINETICS EXPERIMENT | 79 |

LIST OF TABLES

| | |
|--|----|
| Table 1: Characteristics of uranium isotopes in natural uranium | 17 |
| Table 2: Trace metal, metalloid, uranium and thorium concentrations in the Beaverlodge Lake..... | 19 |
| Table 3: Final composition of Hutner's medium and P05-medium | 23 |
| Table 4: Overview of the primers for the potential U-biomarker genes and the new housekeeping genes (used for the normalization of the data) with their related function | 26 |
| Table 5: Overview of the primers for the existing housekeeping genes (used for the normalization of the data) with their related function..... | 28 |
| Table 6: Visual scoring of the comet assay | 29 |
| Table 7: Percentage of growth inhibition (%I _r) calculated from the FW of the plants linked to the total area of L. minor plants exposed for 1, 4 or 7 days to U (0, 2 or 10 µM) and/or metal mixtures (M0, M1, M100)..... | 38 |
| Table 8: Overview of potential U-biomarker primers analyzed with the four biomarker criteria | 44 |
| Table 9: Overview of different electrophoresis times and washing steps for the performed comet assay | 45 |
| Table 10: Overview of results from the visual scoring of the comet assay | 46 |
| Table 11: Exact composition of the contamination solution used in the 7 days exposure experiment and the time kinetics experiment | 63 |
| Table 12: Specificity testing of potential housekeeping genes for L. minor based on real-time qPCR (amplification plot and melt curve) and primer efficiencies (%) | 65 |
| Table 13: Specificity testing of potential U-biomarker genes for L. minor based on real-time qPCR (amplification plot and melt curve) and primer efficiencies (%) | 71 |
| Table 14: Growth measurements (total area and number of fronds) of L. minor plants at day 0, 4 and 7 in 7 days exposure experiment..... | 75 |
| Table 15: Growth measurements (total area and number of fronds) of L. minor plants (sampled at day 1) at day 0 and 1 in time kinetics experiment | 79 |
| Table 16: Growth measurements (total area and number of fronds) of L. minor plants (sampled at day 4) at day 0 and 4 in time kinetics experiment | 82 |
| Table 17: Growth measurements (total area and number of fronds) of L. minor plants (sampled at day 7) at day 0, 4 and 7 in time kinetics experiment | 84 |

LIST OF FIGURES

| | |
|--|----|
| Figure 1: L. minor plants with floating ruler of 10 mm..... | 21 |
| Figure 2: Coated slide for comet assay with a total of 12 gels | 28 |
| Figure 3: Gene expression levels for the primer optimization of potential U-biomarker genes for L. minor relative to control condition (U0-M0) after exposure of 7 days to U (0, 2 or 10 μ M) and/or metal mixtures (M0, M1, M10, M100 or M1000) ... | 32 |
| Figure 4: Percentage of growth inhibition (%I _r) calculated from the FW of the plants linked to the total area of L. minor plants exposed for 7 days to U (0, 2 or 10 μ M) and/or metal mixtures (M0, M1, M10, M100 or M1000)..... | 35 |
| Figure 5: Gene expression levels of potential U-biomarker genes for L. minor relative to control condition (U0-M0) after exposure for 7 days to U (0, 2 or 10 μ M) and/or metal mixtures (M0, M1, M10 or M100) | 36 |
| Figure 6: Gene expression levels of potential U-biomarker primer 33 for L. minor relative to control condition (U0-M0) after exposure for 1, 4 or 7 days to U (0, 2 or 10 μ M) and/or metal mixtures (M0, M1, M100) | 39 |
| Figure 7: Gene expression levels of potential U-biomarker primer 44 for L. minor relative to control condition (U0-M0) after exposure for 1, 4 or 7 days to U (0, 2 or 10 μ M) and/or metal mixtures (M0, M1, M100) | 40 |
| Figure 8: Gene expression levels of potential U-biomarker primer 46 for L. minor relative to control condition (U0-M0) after exposure for 1, 4 or 7 days to U (0, 2 or 10 μ M) and/or metal mixtures (M0, M1, M100) | 41 |
| Figure 9: Gene expression levels of potential U-biomarker primer 48 for L. minor relative to control condition (U0-M0) after exposure for 1, 4 or 7 days to U (0, 2 or 10 μ M) and/or metal mixtures (M0, M1, M100) | 42 |
| Figure 10: Gene expression levels of potential U-biomarker primer 52 for L. minor relative to control condition (U0-M0) after exposure for 1, 4 or 7 days to U (0, 2 or 10 μ M) and/or metal mixtures (M0, M1, M100) | 43 |
| Figure 11: Visual comets of control condition at 15 min of electrophoresis | 45 |
| Figure 12: Results of comet assay | 46 |

ABBREVIATIONS

| | |
|--------------------|--|
| %GC | percentage of guanine and cytosine |
| %I _r | percentage growth inhibition |
| μ | average specific growth rate |
| <i>A. thaliana</i> | <i>Arabidopsis thaliana</i> |
| BIS | Biosphere Impact Studies |
| Ca | calcium |
| Cd | cadmium |
| cDNA | complementary DNA or copy DNA |
| Cu | copper |
| DNA | deoxyribonucleic acid |
| EC _x | effect concentration where the growth is inhibited with X% |
| FW | fresh weight |
| <i>L. minor</i> | <i>Lemna minor</i> |
| M0 | no addition of metals |
| M1 | metal concentration as it occurs in the Beaverlodge Lake (0.0157 μM Cu, 0.017 μM Ni, 0.0765 μM Zn and 0.0145 μM Pb) |
| M10 | tenfold metal concentration of M1 |
| M100 | hundredfold metal concentration of M1 |
| M1000 | thousandfold metal concentration of M1 |
| MES | 2-(<i>N</i> -morpholino)ethanesulfonic acid |
| Mg | magnesium |
| MMS | methyl methanesulfonate |
| mRNA | messenger RNA |
| Na | sodium |
| NCBI | National Center for Biotechnology Information |
| Ni | nickel |
| NMPA | Normal Melting Point Agarose |
| OECD | Organisation for Economic Co-operation and Development |
| Pb | lead |
| PBS | Phosphate Buffer Solution |
| PCR | polymerase chain reaction |
| qPCR | quantitative polymerase chain reaction |
| RNA | ribonucleic acid |
| SE | standard error |
| Sr | strontium |
| t _d | doubling time |
| U | uranium |
| U0 | condition with 0 μM U |
| U10 | condition with 10 μM U |
| U2 | condition with 2 μM U |
| Zn | zinc |
| β | beta |
| γ | gamma |

ABSTRACT

Historical uranium (U) mining operations resulted in the contamination of large areas, including surface waters. The effects of U on the freshwater plant *Lemna minor* have been investigated, although they were mainly studied on individual and cellular level. Sequencing of the *L. minor* genome and transcriptome gives possibilities to study the effects on molecular level. The aim of this study is to select and validate potential U-biomarker genes that can possibly predict the effects of U exposure on population level as well as on ecosystem level. Since organisms growing in contaminated waters are not only exposed to U, but also to multiple co-contaminants, it is necessary to find biomarkers that react specifically to U and not to the co-contaminants.

L. minor plants were exposed to 0, 2 or 10 μM U in combination with an environmentally relevant metal mixture consisting of Cu, Ni, Zn and Pb. Plants were exposed during 1, 4 or 7 days. Expression levels of 10 primers for potential U-biomarkers genes were tested using real-time qPCR. A comet assay on *L. minor* was optimized as a second potential biomarker for DNA damage.

The earlier discovered primer 33 could be validated as biomarker for U exposure. In addition, based on gene expression analysis two new U-biomarkers were selected within this project, i.e. primer 48 and primer 52, since their related genes reacted specifically to U already after 4 days of exposure and to the lowest tested U concentration (2 μM). However, further investigations and validation is necessary.

ABSTRACT IN DUTCH

Historische ontginning van uranium (U) heeft geleid tot contaminatie van grote gebieden, zoals oppervlaktewaters. Effecten van U op de zoetwaterplant *Lemna minor* werden voornamelijk op individueel en cellulair niveau bestudeerd. Door de recente sequencerij van het *L. minor* genoom en transcriptoom kunnen de effecten nu ook op moleculair niveau bestudeerd worden. Het doel van deze studie is om potentiële U-biomarker genen te selecteren en te valideren, die mogelijk de effecten van U blootstelling op populatie- en ecosysteemniveau kunnen voorspellen. Omdat organismen in verontreinigde wateren niet alleen zijn blootgesteld aan U maar ook aan meerdere co-contaminanten, is het noodzakelijk om biomarkers te vinden die specifiek op U reageren en niet op deze co-contaminanten.

L. minor planten werden blootgesteld aan 0, 2 of 10 μM U in combinatie met een relevant metaalmengsel bestaande uit Cu, Ni, Zn en Pb. Planten werden blootgesteld gedurende 1, 4 of 7 dagen. Expressieniveaus van 10 primers voor potentiële U-biomarker genen werden getest m.b.v. real-time qPCR. Een comet assay op *L. minor* werd geoptimaliseerd als tweede potentiële biomarker.

De eerder ontdekte primer 33 werd gevalideerd als biomarker voor U blootstelling. Daarnaast werden ook twee nieuwe U-biomarker genen geselecteerd in dit project op basis van genexpressieanalyse, nl. primer 48 en primer 52. Dit omdat hun verwante genen specifiek reageerden op U al na 4 dagen blootstelling en op de laagste geteste U concentratie (2 μM). Meer onderzoek en validatie is echter nodig.

1 INTRODUCTION

The SCK•CEN is the Belgian Nuclear Research Centre that was established in 1952. One of its goals is to perform research in the nuclear field in order to improve the protection of humans and the environment against ionizing radiation, radionuclides and heavy metals [1].

This thesis is performed in the institute Environment, Health and Safety, and more specifically in the research group Biosphere Impact Studies (BIS). This group has three interlinked research lines. First, there is the understanding of processes and mechanisms for transfer of radionuclides in the biosphere. Process-based models are developed to assess radionuclide dispersion or cycling within different environmental scenarios. Secondly, the group studies the mechanisms behind effects induced in plants following exposure to radiation or radionuclides. Finally, the group develops, improves and applies models for assessing radiological impact on man and environment. This thesis is a part of the second research line. Potential uranium-biomarker genes are selected and validated for the uranium (U) toxicity to *Lemna minor* (*L. minor*), a freshwater plant, in an environmentally relevant metal mixture consisting of copper (Cu), nickel (Ni), zinc (Zn) and lead (Pb).

In the past, several studies have been accomplished with *L. minor* and exposure to radiation/radionuclides at SCK•CEN: 1) uptake and effects of U on *L. minor* in combination with different concentrations calcium (Ca), magnesium (Mg), sodium (Na) and pH [2]; 2) U-cadmium (Cd) mixture experiments on *L. minor*: uptake and growth [3], [4]; 3) recovery of *L. minor* after exposure to gamma (γ) radiation, U or Cd [5]; 4) long-term low dose radiation experiments [6] and 5) effects of γ , strontium (Sr) or U on *L. minor* and ribonucleic acid (RNA)-sequencing [7], [8], [9].

Previously, the effects on *L. minor* plants were only studied on individual and cellular level. The growth inhibition was measured whereby the number and the surface of the leaves, called fronds, were determined. The sequencing and assembly of the *L. minor* genome and the RNA-sequencing of the whole *L. minor* transcriptome by Van Hoeck et al. in 2015 [7] gives new opportunities to study molecular changes of exposed *L. minor* plants. The availability of the transcriptome opens new opportunities for the discovery of biomarkers that can possibly predict the effects of exposure on population level as well as on ecosystem level. In addition, previous experiments mainly studied the effects of single element toxicity on *L. minor*, but it is known that heavy metals can interact with each other. Metals can act independently or have additive, synergistic or antagonistic effects [10]. Therefore, there is an increasing need to study these combined effects. The environmentally relevant metal mixture is based on concentrations that were measured in the Beaverlodge Lake in Canada [11]. As a result of the established U mines local surface waters were contaminated with U and some co-contaminants. The most important co-contaminants (Cu, Ni, Zn and Pb) were selected and used in this research. Prior to this project, in one study the single dose response curves of the different metals of the Beaverlodge Lake were investigated, and a first metal mixture experiment was carried out leading to the discovery of a potential U-biomarker gene (primer 33) [12]. This gene needs to be further investigated and validated by new experiments.

The aim of this research is to validate the potential U-biomarker gene (primer 33) and to select and validate other potential U-biomarker genes for the freshwater plant *L. minor* when they are exposed to U in combination with an environmentally relevant metal mixture. The selection is done based on the available RNA-sequencing data [7]. The first goal is to find genes that specifically react to U and genes that are up- or downregulated,

but only when U is present in the mixture. These genes can then in the future be used as potential U-biomarkers. This will be done by determining the effects on the gene expression levels of selected genes in a 7 days exposure experiment of *L. minor* plants to U in combination with the co-contaminants. The second goal is to find fast-reacting biomarkers in a time kinetics experiment. Hereby the gene expression of the selected genes is followed in function of time by harvesting plants after 1, 4 and 7 days of exposure.

Besides changes at gene expression level, deoxyribonucleic acid (DNA) damage can possibly be used as biomarker for U exposure. Therefore, a comet assay on *L. minor* plants is performed. Since it is the first time that this technique will be performed on *L. minor* within the BIS group, the aim is to test the feasibility of this technique on *L. minor* plants and to optimize some of the parameters. The optimization of the comet assay is performed with plants exposed to methyl methanesulfonate (MMS, i.e. an inducer of DNA damage).

2 ENVIRONMENTAL POLLUTION

All living organisms are exposed to environmental pollution coming from, for example, disposal of industrial effluents, modern agricultural practices, mining and milling... [13]. This can not only affect the human population, but also animals and plants resulting in an increasing worldwide health concern [14]. Heavy metals are important contributors to the pollution. Although they are mostly naturally occurring elements, when they are released in high concentrations changes on cellular and individual level can occur since they are non-biodegradable, potentially toxic to biota and can bio-accumulate [15], [16]. In addition, also radionuclides are an important source of contamination. Radionuclides can be naturally occurring (e.g. U-238, Co-60) or artificially produced (e.g. Cs-137), but both are present in all ecosystems by nature, accident or human activities [17]. In addition historical U mining operations resulted in the contamination of large areas, including surface waters. Therefore there is a need to study the effects of U and heavy metals on plants to preclude major problems to the environment. The selection of potential biomarkers can help us with the prediction of these effects.

2.1 Uranium

Uranium is a naturally occurring radionuclide as well as a heavy metal that is present in aquatic and terrestrial environments. The average concentration of dissolved U in freshwater is 0.03-2.1 $\mu\text{g.L}^{-1}$ [18], [19], [20]. Due to anthropogenic activities such as U mining and milling, processing of U-containing ores and processing of U as fuel for nuclear reactors, high concentrations of U can be found that can reach up to 0.45 mg.L^{-1} which lead to increased toxicological and ecotoxicological concerns [11], [20], [21].

Natural U consist of three isotopes: U-238 (99.27%), U-235 (0.72%) and U-234 (0.0055%) [22], [23]. All of these U isotopes have different radioactive properties, but the chemical properties are the same. The radioactivity depends on the half-life, which is given in Table 1 along with the relative mass and specific activity of the three U isotopes [21], [24], [25]. U is characterized with a higher chemical toxicity than radiotoxicity because the decay half-life is very long [21], [23], [24], [26].

Table 1: Characteristics of uranium isotopes in natural uranium [24, p. 95]

| Isotope | Half-life (years) | Relative mass (%) | Specific activity (Bq.g^{-1}) |
|------------------|--------------------|-------------------|--|
| ^{238}U | 4.47×10^9 | 99.3 | 12455 |
| ^{235}U | 7.04×10^8 | 0.72 | 80,011 |
| ^{234}U | 2.46×10^5 | 0.0055 | 231×10^6 |

The chemical toxicity depends on the U species which can be divided into three predominant species: uranyl cation (UO_2^{2+}), uranyl carbonates (UO_2CO_3 , $\text{UO}_2[\text{CO}_3]_2^-$), and uranyl hydroxides (UO_2OH^+ , $[\text{UO}_2]_3[\text{OH}]_7^-$) [18], [27]. The U speciation is controlled by the pH value, redox potential, ionic strength, and the presence of complexing agents [18], [20], [21]. The uranyl cation (UO_2^{2+}) is the most common form in the environment and it can form complexes with carbonate, phosphate or sulfate ions [23], [24]. It is suggested that U toxicity is mainly caused by UO_2^{2+} due to a number of factors: 1) UO_2^{2+} can replace Ca and Mg ions, which results in structural changes in cell membranes, enzyme inactivation, and damage to RNA and DNA; 2) it can interact with phosphate groups, DNA and it can also induce membrane damage resulting in more permeable membranes; 3) it

can interact with carboxylic groups; 4) it can bind to DNA causing mutations in cells [18], [27], [28].

U is, along with other heavy metals and radiation, a genotoxic agent that changes the balance in chromatin metabolism which results in loss of genomic stability and lower rate of cell division. Genotoxicity can be described as the possibility to destructively effect the genetic material (DNA or RNA) of a cell which can lead to mutations [29]. U can induce DNA damage directly or indirectly by the production of free radicals or reactive oxygen species or by metabolic activation [29], [30], [31], [32]. DNA damage can occur as single-strand breaks, double-strand breaks, alkali-labile sites, incomplete excision repair sites, DNA crosslinks... depending on the concentration or oxidation state of the metal or the exposure time [29], [33], [34]. Studies with the zebrafish *Danio rerio* proved that U is genotoxic and can induce oxidative stress [35], [36]. U also affects the expression of genes of e.g. the detoxification and DNA repair [28].

Humans are exposed to U through food, water and air and the daily intake is approximately $0.037 \mu\text{g}\cdot\text{kg}^{-1}$ of body weight [37], [38]. In general the effects of U in humans are mainly caused by the chemical toxicity (chemotoxic) [38], [39]. U can enter the body in different ways and has also multiple targets. There are targets with a low risk e.g. lymph nodes, digestive tract, muscles, nervous tissue, liver, spleen and blood; medium risk e.g. lungs and gonads and high risk e.g. bones and kidneys. The kidneys have the most chemotoxic risk whereby U binds on biological molecules [40].

Plants are easily exposed to high concentrations of U [41], [42]. Previous experiments with plants showed that U has important effects on growth, nutrient uptake and oxidative stress related responses in plants [43], [44], [45], [46]. But the mechanisms for the uptake and tolerance of U have not yet been fully understood [47]. However experiments with *Arabidopsis thaliana* (*A. thaliana*) have shown that the concentration of U is higher in roots than in shoots due to the adsorption on the cell wall reducing the root-to-shoot translocation [18], [44], [47], [48]. It is not clear yet if experiments with aquatic plants such as *L. minor* give the same results, as they are exposed differently to U (see paragraph 3.1) [49].

2.2 Environmentally relevant metal mixture

In Uranium City, located in northern Saskatchewan Canada, historical U mining operations produced a total of 25939 tonnes of U_3O_8 between 1953 and 1982. This resulted in the contamination of local surface waters like lakes and streams, including the Beaverlodge Lake. Organisms in this area are not only exposed to U, but also multiple co-contaminants (Table 2). This results in a need to determine effects from contaminants in real environmental conditions, including metal mixtures [11], [50], [51]. The co-contaminants of interest from the Beaverlodge Lake are Cu, Ni, Zn and Pb. They are released by natural or anthropogenic activities and can be assimilated by aquatic organisms via different routes possibly leading to toxic effects [52], [53], [54], [55].

Table 2: Trace metal, metalloid, uranium and thorium concentrations in the Beaverlodge Lake [11, p. 102]

| Contaminants | Concentration ($\mu\text{g}\cdot\text{dm}^{-3}$) |
|--------------|--|
| As | 1.8 |
| Cu | 1 |
| Mn | nm |
| Fe | 65 |
| Ni | 1 |
| Zn | 5 |
| Se | 4.8 |
| Mo | nm |
| Ba | 0.56 |
| Pb | 3 |
| U | 483.6 |
| Th | 79000 |

*nm = not measured

Cu is an essential micronutrient for all organisms, including plants, and is involved in various physiological processes such as photosynthesis, respiration, cell wall metabolism and hormone signaling [54], [56], [57]. In addition, it is a cofactor in various enzymes [57], [58]. However, an excess of Cu can lead to toxic effects e.g. by forming free radicals which can cause oxidative stress [52]. Cu in freshwater is normally found in concentrations of $0.44 \mu\text{g}\cdot\text{L}^{-1}$, but due to anthropogenic activities high concentrations of Cu were found, which are not only toxic to plants but also to human and animals. The concentration in the plants depends not only on the plant species, but also on the growth stage [52], [56], [57], [59].

Ni is found at concentrations from $1\text{-}3 \mu\text{g}\cdot\text{L}^{-1}$ in freshwaters and can reach up to $10\text{-}15 \mu\text{g}\cdot\text{L}^{-1}$ in polluted areas [54], [60], [61]. It is an essential micronutrient for plants, e.g. it is an important element of the enzyme urease and thus essential for plants which produce this enzyme [58]. Since Ni is a micronutrient, a Ni deficiency is rare but an excess of Ni is more common leading to toxic effects. Depending on the plant species, growth stage, cultivation conditions, Ni concentration and exposure time, different effects are ascertained [60]. For example, Ni can cause inhibition of growth, chlorosis and necrosis in plants [62].

Zn is a non-ferrous metal that in freshwaters is found in concentrations of $2.8 \mu\text{g}\cdot\text{L}^{-1}$ [63]. It is an essential micronutrient for all biota [53], [64]. It has an important role in the physiology of the cell and for the structure of proteins [54]. It is also necessary for enzymes that have a role in carbohydrate and protein synthesis, gene regulation, structure and integrity of biomembranes and protection of cells from damage due to free radicals [58]. It is also known that Zn reduces metal toxicity (e.g. Cu and Cd) which improves the plant growth [65]. But toxic Zn concentrations no longer have the protective function and can cause ecotoxicological effects on plants such as reduced biomass, leaf chlorosis and root growth inhibition [64], [65], [66].

Pb is a toxic element that is naturally present in small quantities in the earth's crust. Anthropogenic activities increase the concentration of Pb in the environment, which results in elevated Pb concentrations in surface and groundwater's [67], [68], [69]. In freshwater the concentration of Pb can reach up to $0.18\text{-}1 \mu\text{g}\cdot\text{L}^{-1}$ [70]. Pb occurs as inorganic Pb(II)

and is one of the few metals that has no biological function [69], [71], [72]. It is known that Pb(II) can disrupt biological systems by modifying molecular interactions, cell signaling and cellular function [68], [71]. Plants mainly absorb Pb(II) in their roots, translocation to the shoots is not very common. Pb(II) can accumulate and critical levels will affect the plant itself by a decrease of biomass or will eventually kill it [62], [73], [74].

Although the toxic effects of the single metals have been investigated before, the environment is contaminated with more than one heavy metal. In addition, the concentration of the metals is not constant. When the concentration rises above the tolerated concentration, toxicity effects appears and changes can be induced [54], [56], [75]. This is not caused by the individual metals itself, but by those mixtures of metals that can have joint effects. The observed effects with co-contaminants can diverge from the results of single-stressor experiments. Metals can have additive, synergistic or antagonistic effects [76] or act independently, depending on the concentration, temperature, pH and light intensity [10]. For example, Vellinger et al. (2012) [77] found an antagonistic metal interaction of arsenate and Cd on *Gammarus pulex*. Also Charles et al. (2006) [78] found an antagonistic effect, but between U and Cu in *Lemna aequinoctialis*. The metal effects are also dependent on the relationship of the metals to other mineral elements. Therefore more research is needed to study the joint effects of the metal mixtures [75], [79], [80].

2.3 *Lemna minor*

The freshwater plant used in this study is *L. minor* (Figure 1). It is an aquatic plant that has an important role in aquatic ecosystems. It is a free-floating freshwater macrophyte that belongs to the subfamily of duckweeds (*Lemnaceae*). It is a relatively small vascular plant from a few mm up to 1 cm [21], [81]. These plants consists of one, two or three leaves, called fronds, which have a single root. The roots are approximately 1-2 cm long, whereas the fronds are oval shaped with a length of 0.6-5 mm thick, light green in color with three nerves and a small air pocket to advance floating. For their growth the conditions should be as ideal as possible: high level of nutrients, pH between 5 and 9 (optimum 6.5-7.5), and temperature between 6 and 33°C [81]. They have a high growth rate via asexual reproduction. On average, one adult plant splits into two new plants every 2.5 days. This results in identical clones [7], [81], [82]. As a result of the genetically identical clones potential effects due to genetic variability can be eliminated [7].

Plants are essential to a healthy ecosystem. The duckweed species have an ecological significance as the primary producers of food for waterfowl, fish and small organisms, and oxygen [7], [82]. *L. minor* plants are sensitive towards different kinds of stress (e.g. heavy metals) and are able to absorb and accumulate high amounts of it in their biomass [52], [56], [82], [83]. Due to their easy growth in controlled and sterile laboratory conditions *L. minor* is often used as a model test organism for environmental studies and fundamental plant research [7], [82]. The Organisation for Economic Co-operation and Development (OECD) describes a growth inhibition test for *L. minor* (guideline 221) to test the toxicity of pollutants in existing freshwaters [21], [52], [56], [84]. Toxicity of compounds is expressed in EC_x, the effect concentration where the growth is inhibited with X% [81].




Figure 1: *L. minor* plants with floating ruler of 10 mm

2.4 Biomarkers

The interest for biomarkers is still growing to determine the toxic effects on the environment. They can be used as early warning systems for organisms that are exposed to single or complex mixtures of chemicals resulting in a changed biological response [81], [85], [86]. There are two different types: biomarkers of exposure and biomarkers of effect. Biomarkers of exposure are used as functional measure of exposure to toxicants, whereby the expression is on a suborganism level. Biomarkers of effect, also called bioindicators, describes changes in biochemical, physiological or ecological parameters that are a consequence of exposure to a toxicant [87], [88], [89], [90]. Terrestrial and aquatic plants can be early warning systems to measure the toxicity in natural freshwaters. When they are exposed to toxicants, the first effects are perceived at molecular and biochemical level and afterwards on level of the organism [81], [85], [87].

There are multiple potential biomarkers of exposure of which gene expression is one of them. Changes in gene expression give the first effects of exposure and are therefore highly sensitive information [16], [91], [92]. It can be used to study the mechanisms behind exposure and form a 'genetic signature' of the specific exposure pattern. Exposed plants can up- or downregulate specific genes dependent on the exposure [87]. The most commonly used techniques for gene expression analysis are RNA sequencing, Northern blot, microarray analysis and polymerase chain reaction (PCR). This last technique will be used in this project. Multiple genes at messenger RNA (mRNA) level can be simultaneously studied and potential molecular mechanisms and biomarkers can be identified. But there are also certain disadvantages such as variation of genes due to intern variability (physiologic state, cell types, environmental conditions...) [87], [92].

Another potential biomarker of exposure is DNA damage. As described above, genotoxic agents such as heavy metals or radiation can cause DNA damage. The genotoxicity of single genotoxic agents and mixtures can be evaluated by single-cell gel electrophoresis, also called "comet assay", whereby the effects of genotoxic agents are studied on individual cells [93]. Cells are embedded in agarose resulting in immobilized DNA. Subsequently an electrophoresis is performed whereby the DNA is migrated and the nuclei are fixated and afterwards indicated with a fluorescent staining [86]. In the past this technique was mainly

used with animal cells, but the use on plants gives new opportunities in environmental studies [93], [94], although using plants has an important disadvantage compared to the use of animal cells. The cellulose cell wall forms a barrier to prevent the release of DNA. By chopping the plant tissues with a razor blade, more optimal conditions are created for the release of the DNA [95]. The neutral version of the comet assay was first tested on plants, whereby the electrophoresis is performed under neutral conditions. However, the sensitivity decreased with the neutral version [95]. Thereby the alkaline version was created on *Vicia faba* to quantitatively measure DNA damage [96], [97], [98]. Hereby the treatment and electrophoresis is fulfilled under alkaline conditions which results in better visible comet tails and higher sensitivity [95]. This alkaline version of the comet assay has also some other advantages such as short analysis time and it is an inexpensive, non-specific method [31]. But, as described above, due to the plants cell wall and the absence of free cells difficulties arise with performing the comet assay on plants. Also the comet assay is known to be a technique that is difficult to standardize, which results in a lower reliability [96].

3 METHODS AND MATERIALS

In this chapter the methods and materials are given starting with the *L. minor* growth and exposure to U and the metal mixture. Subsequently the calculation of the specific growth rate and growth inhibition, the primer development and optimization and the measurements at molecular level are described. There are two types of measurements on molecular level: gene expression and the comet assay. At last the used statistical analysis is given.

3.1 *L. minor* growth and exposure to uranium and metal mixture

L. minor plants were cultivated in sterile Erlenmeyer flasks of 250 ml that consists of 100 ml sterile growth medium, also called Hutner's medium, and were closed with hydrophobic cotton and aluminum foil. The composition of the Hutner's medium is given in Table 3. Every 10 to 12 days the stock cultures were transferred to new Erlenmeyer flasks to maintain the plant culture. Therefore, three *L. minor* plants with 3 or 4 fronds were transferred with a sterile inoculation loop to a new sterile Erlenmeyer flask. All operations were executed in sterile conditions. According to the OECD guidelines the stock culture needs to be grown at 24°C with continuous light. But due to growth problems of the plants, they were grown under a day and night program (14h/10h) at respectively 24/18°C with a light intensity of $\pm 220 \mu\text{mol}\cdot\text{s}^{-1}\cdot\text{m}^{-2}$. In addition, non-sterile *L. minor* plants were used. All the subsequent proceedings were performed under sterile conditions as described earlier.

Table 3: Final composition of Hutner's medium and P05-medium

| | Hutner's medium (mg.L ⁻¹) | P05-medium (mg.L ⁻¹) |
|--|---------------------------------------|----------------------------------|
| Macronutrients | | |
| KNO ₃ | 0.3 | 888.8 |
| Ca(NO ₃) ₂ ·4H ₂ O | 0.72 | 944 |
| MgSO ₄ ·7H ₂ O | 0.0738 | 500 |
| KH ₂ PO ₄ | 0.0414 | 503.2 |
| Na-EDTA | 0.0029 | 9 |
| Ferric acid | 0.001 | / |
| Tartaric acid | / | 3 |
| Micronutrients | | |
| H ₃ BO ₃ | 0.001 | 1.860 |
| ZnSO ₄ ·7H ₂ O | 0.001 | 0.220 |
| Na ₂ MoO ₄ ·2H ₂ O | 0.0001 | 0.120 |
| CuSO ₄ ·5H ₂ O | 0.00003 | 0.080 |
| MnCl ₂ ·4H ₂ O | / | 3.620 |
| FeCl ₃ ·6H ₂ O | / | 5.400 |
| MnSO ₄ ·H ₂ O | 0.0001 | / |

In total two experiments were performed which followed the standard 7-day growth inhibition test of the OECD using *L. minor*, with some modifications as described by Horemans et al. [99] and the alterations as described above. Six days before the experiments, a preculture of five *L. minor* plants with 3 or 4 fronds was set up in a sterile Erlenmeyer flasks of 250 ml that consists of 100 ml Hutner's medium.

The first experiment was performed with *L. minor* exposed in quadruplicate for 7 days to a metal mixture in combination with U. Five metal concentrations were used: M0, M1, M10, M100 and M1000. The metal concentration M1 is the concentration as it occurs in nature (0.0157 μM Cu, 0.017 μM Ni, 0.0765 μM Zn and 0.0145 μM Pb). M10, M100 and M1000 are respectively a tenfold, hundredfold and thousandfold concentrations and M0 is no addition of those metals. For U three concentrations were selected: 0, 2 and 10 μM , whereby 2 μM is the concentration occurring in the Beaverlodge Lake.

For the contamination a M10000 stock solution of Cu, Ni, Zn and Pb was prepared from separated 100 mM stock solutions of the metals ($\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$ (Sigma-Aldrich), $\text{NiSO}_4 \cdot 6\text{H}_2\text{O}$ (Thermo Fisher Scientific), $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$ (Honeywell Fluka) and $\text{Pb}(\text{NO}_3)_2$ (Sigma-Aldrich)) and filter-sterilized with a 0.2 μm HT Tuffryn Membrane (Pall Corporation). An U stock solution of 10 mM was prepared from $\text{UO}_2(\text{NO}_3)_2 \cdot 6\text{H}_2\text{O}$ (SPI-chem) and filter-sterilized with a 0.2 μm HT Tuffryn Membrane (Pall Corporation). During the exposure, P05-medium (Table 3) was used that is low in phosphate concentration to limit U precipitation [99]. Sterile 250 ml polycarbonate pots were filled with the contamination solution consisting of 0, 0.02 or 0.1 ml U stock solution and 0, 0.01, 0.1, 1 or 10 ml M10000 stock solution depending on the condition and afterwards filled with P05-medium until a total volume of 100 ml (exact composition of each condition is given in Appendix A). At each condition 1 ml of a filter-sterilized (0.2 μm HT Tuffryn Membrane (Pall Corporation)) 0.5 M 2-(*N*-morpholino)ethanesulfonic acid (MES) (Sigma-Aldrich) buffer was added and the pH of each solution was adjusted to 5 with filter-sterilized (0.2 μm HT Tuffryn Membrane (Pall Corporation)) 1 M HCl (VWR) or 1 M NaOH (Merck) to create the same experimental conditions. All *L. minor* plants of the preculture were transferred into a large polycarbonate pot of 1 L and afterwards three plants with 3 or 4 fronds were selected and transferred into the 250 ml polycarbonate pots filled with the P05-medium, U and/or metal contamination solution and MES buffer. Also a piece of sterile, floating ruler was added in each pot for later measurements. Pictures were taken at day 0, 4 and 7 for determinations of the number and surface of fronds (paragraph 3.2). After 7 days, plants were harvested, put onto an absorbent paper to remove any present medium and fresh weight (FW) was determined. Samples from 20-50 mg were snap frozen in liquid nitrogen for RNA extractions.

The second experiment was a time kinetics experiment whereby *L. minor* plants were exposed to three concentrations of the metal mixture (M0, M1, M100) in combination with the three U concentrations (0, 2 and 10 μM). The same experimental set-up as described above was used. Pictures were taken at day 0, 1, 4 and 7 for determinations of the number and surface of fronds (paragraph 3.2). After 1, 4 and 7 days, plants were harvested, FW was determined and samples of 20-50 mg for RNA extractions were frozen in liquid nitrogen to see how the gene expression of the possible biomarkers changes over time. The *L. minor* plants for day 1 of the harvest were exposed in sevenfold, whereas the plants for day 4 and 7 of the harvest were exposed in quadruplicate.

3.2 Specific growth rate and growth inhibition

Three measurements on cellular level were performed: FW (see paragraph 3.1), number of fronds and surface of fronds.

For both experiments pictures were taken to determine the number and surface of fronds at day 0, 4 and 7 (for the 7 days exposure experiment) and at day 0, 1, 4 and 7 (for the time kinetics experiment). The pictures were analyzed using the open source image processing program ImageJ. The total area was determined and the number of fronds was

also counted. The FW of the plants on each time point could be calculated based on the ratio FW-total area on day 0. The total area, FW or number of fronds was used to determine the average specific growth rate (μ) with the following formula [84]:

$$\mu_{i-j} = \frac{\ln(N_j) - \ln(N_i)}{t_j - t_i} \quad (1)$$

with μ_{i-j} as the average specific growth rate from time i to j , N_i the number of fronds observed, frond area or FW at time i , N_j the number of fronds observed, frond area or FW at time j , t_i the time point for the start of the period and t_j the time point for the end of the period.

With the average specific growth rate of the control and the contaminated conditions, the percentage of growth inhibition ($\%I_r$) could be calculated as [84]:

$$\%I_r = \frac{(\mu_c - \mu_T)}{\mu_c} \cdot 100 \quad (2)$$

with μ_c the mean value for μ in the control condition and μ_T the mean value for μ in the contaminated condition.

Also the doubling time (t_d) was measured for the validation of the test. The t_d of frond number in the control condition must be less than 2.5 days (or the average specific growth rate must be higher than 0.275 d^{-1}) and could be calculated from the average specific growth rate with the following formula [84]:

$$t_d = \frac{\ln(2)}{\mu} \quad (3)$$

The FW, average specific growth rate (μ) and percentage growth inhibition ($\%I_r$) were used as endpoints to compare the various conditions.

3.3 Primer development and optimization

Van Hoeck et al. (2015) [7] already studied the gene expression levels of *L. minor* plants exposed to U, beta (β) and γ radiation. This expression data was used to find potential U-biomarker genes that react specifically to U. Genes were selected based on their changed expression level for U with no changes after exposure to β and γ radiation. Genes with an increasing or decreasing fold change in function of increasing U concentration were selected for potential up- or downregulated U-biomarker genes. The mRNA of the selected genes along with the whole *L. minor* transcriptome was used for primer development with the primer-BLAST tool of National Center for Biotechnology Information (NCBI). Preprogrammed settings of NCBI were used, only the PCR product size was changed to 70-150 bp, maximum melt temperature difference was set on 2°C and targets were ignored that had four or more mismatches to the primer. Subsequently the properties and the possibility to form self-dimers of the potential primers were tested with the Oligo Analysis Tool (Eurofins Genomics). Only primers with a percentage of guanine and cytosine ($\%GC$) between 50 and 60% and a melting temperature around 60°C were selected. The maximum annealing score, the number of overlay between two identical primers, should be maximum 14 for a potential good primer. Finally, a PCR check was executed with the Oligo Analysis Tool (Eurofins Genomics) to see if the forward and reverse primer can form cross-dimers during the PCR reaction. The formation of cross-dimers was the result of a homologous forward and reverse primer. If all the requirements were approved the primers

were ordered and further tested and validated within this project (paragraph 3.4.1). The developed primer pairs were given a number. This number (i.e. Primer "X") was used to describe the potential U-biomarker genes where the primers were linked to (see Table 4). Due to confidentiality of this thesis no primer sequences and gene numbers are given.

In addition, new housekeeping genes to improve the normalization of the gene expression data (paragraph 3.4.1) were selected from the RNA sequencing data [7]. Hereby genes with a constant fold change in function of all exposure conditions were selected. This mRNA of the new potential housekeeping genes along with the whole *L. minor* transcriptome was used for primer development as described above. The primers for the housekeeping genes are given in Table 4.

Table 4: Overview of the primers for the potential U-biomarker genes and the new housekeeping genes (used for the normalization of the data) with their related function

| Primer number | Function |
|-------------------------------|-------------------------------------|
| U-BIOMARKER GENES | |
| Primer 33 | CYP76C2 Cytochrome P450 76C2 |
| Primer 41 | Negative regulation of growth |
| Primer 42 | Negative regulation of growth |
| Primer 43 | Negative regulation of growth |
| Primer 44 | Negative regulation of growth |
| Primer 45 | Response to oxidative stress |
| Primer 46 | Response to oxidative stress |
| Primer 47 | Response to oxidative stress |
| Primer 48 | Response to oxidative stress |
| Primer 49 | Metal ion transport |
| Primer 50 | Metal ion transport |
| Primer 51 | Response to auxin |
| Primer 52 | Response to auxin |
| NEW HOUSEKEEPING GENES | |
| Primer 53 | PAL |
| Primer 54 | PAL |
| Primer 55 | Cytochrome |
| Primer 56 | Cytochrome |
| Primer 57 | Fatty acid biosynthetic process |
| Primer 58 | Fatty acid biosynthetic process |
| Primer 59 | Flavin adenine dinucleotide binding |
| Primer 60 | Flavin adenine dinucleotide binding |
| Primer 61 | Flavin adenine dinucleotide binding |
| Primer 62 | Flavin adenine dinucleotide binding |
| Primer 63 | Ubiquitin protein ligase binding |
| Primer 64 | Ubiquitin protein ligase binding |
| Primer 65 | Cytochrome |
| Primer 66 | Cytochrome |
| Primer 67 | Photoreceptor |
| Primer 68 | Photoreceptor |

3.4 Measurements on molecular level

Two types of measurements on molecular level were performed in this thesis. First, the gene expression was analyzed whereby RNA was extracted from plant samples of the 7 days exposure experiment and time kinetics experiment. Subsequently copy DNA (cDNA or complementary DNA) was made and the real-time qPCR (quantitative polymerase chain reaction) reactions were executed. Secondly, the comet assay was performed whereby *L. minor* plants were exposed to MMS to see if induced DNA damage could be measured.

3.4.1 Gene expression

To analyze the gene expression of selected housekeeping genes and possible U-biomarkers, RNA was extracted. For this, frozen plant samples at -80°C were transferred to liquid nitrogen and then disrupted using the Mixer Mill MM400 (Retsch). At each sample approximately ten Zirconia (Biospec) beads (2.0 mm) were added. Afterwards the samples were placed in pre-cooled Mixer Mill adaptors and grounded for 3.5 minutes at 30 Hz. The RNA was extracted using the RNeasy Plant Mini kit (Qiagen). The RNA content was measured with the Nanodrop ND-1000 (Thermo Fisher Scientific) and all samples were set on the same concentration of 10000 ng per 45 µl. RNA samples were stored at -20°C until further use.

Before synthesis of the cDNA, any present DNA in the RNA samples was removed using the Turbo DNA-free Kit (Invitrogen). Afterwards the RNA concentration was measured again with the Nanodrop ND-1000 (Thermo Fisher Scientific) and all samples were set on the same concentration of 1000 ng per 13 µl. Subsequently the cDNA was synthesized using the PrimeScript RT reagent Kit (Perfect Real Time) (TaKaRa) with standard protocol and the Pqstar Gradient thermocycler (VWR). cDNA samples were diluted with RNase and DNase free water (1:10) for real-time qPCR and can be stored at -20°C.

The qPCR was performed with the Rotor-Gene Q (Qiagen), using SYBR green chemistry and the developed primers of paragraph 3.3. For the qPCR the QIAgility (Qiagen) was used to fill each well of a Rotor-Disc with a total volume of 15 µl containing 11.25 µl qPCR MasterMix (7.5 µl Fast SYBR Green MasterMix (Applied Biosystems), 2.85 µl RNase and DNase free water, 0.45 µl forward primer and 0.45 µl reverse primer) and 3.75 µl cDNA. Subsequently the Rotor-Disc was sealed with the Rotor-Disc Heat Sealer (Qiagen) and placed in the Rotor-Gene Q (Qiagen) for the PCR reaction. During the qPCR, the following program was used: an initiation step at 95°C for 20 s followed by 45 cycles (3 s at 95°C and 30 s at 60°C). A melting curve was generated between 50 and 95°C to ascertain the specificity of the primers.

The amplification efficiency of the primers was tested with a mixed cDNA sample dilution series (1; 1/4; 1/16; 1/64; 1/256; 1/1024). The Rotor-Disc was filled with 11.25 µl qPCR MasterMix and 3.75 µl cDNA. The PCR reaction was performed with the Rotor-Gene Q (Qiagen). With the PCR products, a gel electrophoresis was performed to confirm the desired amplicon. The gel consists of 0.8 g agarose (Roche) and 5 µl GelRed (Biotium) per 100 ml TAE buffer. To load the samples on the gel, 1 µl loading dye was added to 5 µl PCR product. Per gel, 4 µl of this PCR product mixture and 2 µL ladder of 50 and 100 bp (Thermo Scientific) was loaded. The gel electrophoresis was ran for 50 min at 70 V and subsequently visualized with a gel imager (BioRad).

Various developed and optimized primers of the potential U-biomarker genes were tested together with multiple existing and new housekeeping genes (Table 4 and Table 5). The expression stability of the housekeeping genes was tested using GrayNorm [100]. Data

were normalized with a normalization factor from the housekeeping genes. Mean values of the gene expression of the U-biomarker genes was set relative to the control conditions by using the $2^{-\Delta Ct}$ method.

Table 5: Overview of the primers for the existing housekeeping genes (used for the normalization of the data) with their related function

| EXISTING HOUSEKEEPING GENES | |
|------------------------------------|--|
| Primer number | Function |
| Primer 4 | BSL2 Serine/threonine-protein phosphatase BSL2 |
| Primer 9 | SBT3.3 Subtilisin-like protease SBT3.3 |
| Primer 15 | Mitogen-activated protein kinase kinase 1 |
| Primer 18 | TUBB5 Tubulin beta-5 chain |
| Primer 20 | CYP71A25 Cytochrome P450 71A25 |
| Primer 22 | GCP3 Gamma-tubulin complex component 3 |

3.4.2 Comet assay

For the first time a comet assay on *L. minor* plants was performed within the BIS group. The comet assay was performed to optimize some of the parameters. Two conditions were fulfilled: plants exposed to MMS (Sigma-Aldrich) which is an inducer of DNA damage and a control condition. MMS serves as a positive control for DNA damage. In a time period of four hours, the *L. minor* plants were exposed to 5 mM MMS. Afterwards the comet assay handlings were executed.

At least one day before the experiment slides need to be coated with a 1% Normal Melting Point Agarose (NMPA) (Sigma-Aldrich). The NMPA was boiled and homogenated several times. Subsequently the slides were submerged into the NMPA for at least three times. The coated slides were dried overnight.

The comet assay was performed in the dark with only an inactinic red light to avoid the induction of DNA damage through actinic light. The first step was chopping of the plant material. For this, two *L. minor* plants were put into a petri dish and 400 μ l extraction buffer (1/10 of 200 mM Na₂EDTA solution pH 10 (Sigma-Aldrich) and 9/10 of Phosphate Buffer Solution (PBS) pH 7 (consisting of 0.2 g KCl (Sigma-Aldrich), 0.2 g KH₂PO₄ (Merck), 8 g NaCl (Merck) and 1.78 g Na₂HPO₄·2H₂O (Merck) per liter)) was added. Plants were chopped with a razor blade during 30 s. Hereafter, 225 μ l of the nucleus solution was collected and mixed with 150 μ l 2% Low Melting Point Agarose (Sigma-Aldrich). Subsequently, three gels of 10 μ l were pipetted on the coated slide as shown in Figure 2. These operations were repeated until a total of 12 gels were pipetted on the coated slide.




Figure 2: Coated slide for comet assay with a total of 12 gels

The slides were placed into the electrophoresis tank parallel to the border, filled with electrophoresis solution (3 ml of the Na₂EDTA solution and 18 ml of 10 N NaOH (Merck) in a total of 0.5 L). The denaturation took place during 15 min whereby the DNA in the gels






unwinds. Then an electrophoresis was ran at 300 mA and 16-18 V at 4°C. Three electrophoresis times were tested: 5, 10 and 15 min.

After the electrophoresis, the slides were neutralized for 1 min in demineralized water and twice for 5 min with 2 ml of the PBS. The gels were fixated overnight in absolute ethanol (Merck) until they were completely dry.

Slides were stained with 2 ml of a 1/10000 Sybr Gold solution (Invitrogen) after which the slides were washed twice with demineralized water to reduce the amount of background fluorescence. Two washing steps were tested: two washing steps of 5 min and a washing step of 5 min followed by a washing step of 10 min. Comets were visualized using the fluorescence microscope (Nikon Eclipse Ti with Lumencor Spectra X Chroma) with a 20x magnification.

A visual scoring was applied as described by Azqueta et al. (2011) [101] to determine the different levels of DNA damage. A scheme for visual scoring based on 5 recognizable classes of comets (Table 6) was used.

Table 6: Visual scoring of the comet assay [101], [102]

| Class | Description | Picture |
|-------|--|---|
| 0 | Undamaged, no discernible tail |  |
| 1 | Very little damaged |  |
| 2 | Normal damage, nice comet |  |
| 3 | Much damage |  |
| 4 | Almost all DNA in tail, insignificant head |  |

3.5 Statistical analysis

Statistical analyses were performed using the open source software package R (R Project for Statistical Computing, version 3.5.3). Outliers were detected with a Grubbs' test (GraphPad), also called the ESD method (Extreme Studentized Deviate), with a significance level of 0.05.

Data without outliers was used to detect differences between groups using two-way ANOVA. First a Shapiro-Wilk normality test and a Bartlett test were performed to test the normality of the data and homogeneity of the variances, respectively. If the assumption of normality of homogeneity of variances was not fulfilled, transformations of the data were applied where necessary. Four different transformations were performed: $\log(x)$, $(x)^{1/2}$, $1/x$ and e^x . The post-hoc Tukey HSD test was used for multiple comparison of the data.

If the assumptions for a parametric test were not met, a non-parametric Kruskal-Wallis test was performed, followed by a Pairwise Wilcoxon signed-rank test with no correction.

4 RESULTS

In this chapter, the results of this thesis are presented starting with the primer selection, optimization and testing their selectivity. Secondly, the growth and gene expression analyses of the 7 days exposure experiment to a metal mixture in combination with U are described followed by the time kinetics experiment of *L. minor* plants. Finally, the results of the comet assay performed on *L. minor* plants are presented.

4.1 Primer selection and optimization

As described above, the Beaverlodge Lake contains multiple contaminants, with the most important ones being U, Cu, Ni, Zn and Pb. The sequencing of the *L. minor* genome and transcriptome gives the possibility to study changes at the molecular level after exposure to those contaminants. Primers were developed using the primer-BLAST tool of NCBI and further tested with the Oligo Analysis Tool (Eurofins Genomics). Only primers with a %GC between 50 and 60% and a melting temperature around 60°C were selected. Due to the growth problems of the sterile plants, the efficiency of all new primers, 12 potential U-biomarker genes and 16 housekeeping genes (designed in paragraph 3.3) were in first instance tested on available cDNA of a previous U and metal mixture experiment [12]. For each primer the amplification plot and the melting curve are analyzed. A primer that reacts specific to one gene is given by one peak in the melting curve. In addition, the expression of the dilution series is concentration-dependent with the highest expression in the undiluted sample. The efficiencies of all primers is evaluated by setting the threshold in the linear phase of the amplification plot and using the correlated Ct values (Appendix B and C). Primers with an amplification efficiency of at least 90% and approved amplification plot and melting curve were selected and used for further testing.

4.2 Primer selectivity

The primers with a primer efficiency >90% were further tested as potential biomarker for U exposure. This was done with cDNA samples of a previous U and metal mixture experiment (*L. minor* plants exposed for 7 days to U (0, 2 or 10 µM) and/or metal mixtures (M0, M1, M10, M100 or M1000)) [12]. The tested primers were: primer 33, 41, 43, 44, 46, 47, 48, 50, 51 and 52. The expression of the potential U-biomarker genes was normalized with the new and existing housekeeping genes. Gene expression levels relative to the control condition (U0-M0) are given in Figure 3.




Figure 3: Gene expression levels for the primer optimization of potential U-biomarker genes for *L. minor* relative to control condition (U0-M0) after exposure of 7 days to U (0, 2 or 10 μ M) and/or metal mixtures (M0, M1, M10, M100 or M1000). Mean values are given \pm standard error (SE) of at least three biological independent replicates for each condition. Significant differences (two-way ANOVA, p -value <0.05) within the U0 condition are given with small letters, within the U2 condition with capital letters and within the U10 condition with small letters and an apostrophe. Significant differences (two-way ANOVA, p -value <0.05) between the U0 condition and U2/U10 condition are given with an asterisk.




Figure 3 (continued).

In general, there is no significant effect of the different metal mixtures applied within one U concentration, except for the M1000 where significant changes are often observed. This can be explained by the fact that the M1000 conditions are highly concentrated and cause high toxicity to the plants [12]. These conditions cannot be used as early warning systems, which was the initial goal of this thesis. Therefore, the results of the M1000 conditions will not be further discussed in this thesis.

In the absence of U, no significant effects are observed for primer 33 between the different metal concentrations applied. The same trend is present in the U2 and U10 conditions. In addition, primer 33 shows a strong U-dependent response with an upregulation after exposure to 2 (non-significant) and 10 μ M U (significant) in the absence of the metal mixture. At the U concentration of 10 μ M the gene expression is approximately 22 times higher than the control condition. The M0, M1, M10 and M100 of the U10 condition are significant higher than their respective control without U.

For primer 41 no effect of the metal mixture is observed for the different U concentrations applied (0, 2 or 10 μ M). In addition, no effects are present after exposure to U2. There is a significant increase in the gene expression after exposure to U10 as compared to U0 for all metal mixtures applied (except for U10-M1).

Primer 43 and primer 44, both primers for the same gene, show the same trend. No effect of the metal mixture is observed for the different U concentrations applied (0, 2 or 10 μ M).

In addition, no effects are present after exposure to U2. There is a significant increase in the gene expression after exposure to U10 as compared to U0 for all metal mixtures applied.

For primer 46, no significant effects are observed between the different metal mixtures and control plants within one U concentration. An U-dependent up regulation is obtained with a significant effect after exposure to 10 μ M U compared to the plants without U, but the upregulation is lower in comparison with the previous primers. An up regulation of approximately 5 is acquired for the U10 condition, whereas an up regulation of approximately 2 for the U2 condition.

Since primer 47 and primer 48 are both primers for the same gene, the results of both primers are similar. Within one U concentration, no significant effects are present, except for the downregulation in the U2-M100 condition compared to U2-M0 in primer 48. There is an U-dependent downregulation of the expression, whereby the expression level at U10 almost drops to zero.

Primer 50 does not give clear results. In general, no significant results are obtained.

Primer 51 and primer 52 are both primers for the same gene and the results show high similarities. For the different metal mixtures tested, no significant changes are observed within one U concentration. After exposure to U, a significant downregulation is obtained whereby M0 and M1 of the U2 condition and M0, M1, M10 and M100 of the U10 condition are significant lower than their corresponding control.

Based on the results of Figure 3, a selection of four primers is made, besides primer 33, to test in the 7 days exposure experiment (paragraph 4.3) and the time kinetics experiment (paragraph 4.4). Primer 44 and primer 46 as upregulated genes and primer 48 and primer 52 as downregulated genes.

4.3 7 days exposure experiment

During the 7 days exposure experiment *L. minor* plants were exposed to five concentrations of the metal mixture (M0, M1, M10, M100 and M1000) in combination with three U concentrations (0, 2 and 10 μM). The growth of the *L. minor* plants was followed by measuring the total area of the plants and counting the number of fronds (Appendix D). In this thesis the percentage of growth inhibition ($\%I_r$) (Figure 4) is calculated from the FW of the plants linked to the total area.




Figure 4: Percentage of growth inhibition ($\%I_r$) calculated from the FW of the plants linked to the total area of *L. minor* plants exposed for 7 days to U (0, 2 or 10 μM) and/or metal mixtures (M0, M1, M10, M100 or M1000). The percentage of growth inhibition of the control condition (U0-M0) is 0%. Mean values are given \pm standard error (SE) of at least four biological independent replicates for each condition. Significant differences (two-way ANOVA, p -value < 0.05) within the U0 condition are given with small letters, within the U2 condition with capital letters and within the U10 condition with small letters and an apostrophe. Significant differences (two-way ANOVA, p -value < 0.05) between the U0 condition and U2/U10 condition are given with an asterisk.

Exposing plants to M1000 causes $>50\%$ growth inhibition in all conditions, indicating high toxicity to the plants, as mentioned before in paragraph 4.1. These conditions cannot be used as early warning systems, which was the initial goal of this thesis. Therefore, the results of the M1000 conditions will not be further discussed.

In general, no significant differences are observed for the percentage of growth inhibition between the different U exposures. However, in the absence of the metal mixture, growth is reduced with $>20\%$ (non-significant effect). In addition, no significant effect is observed of the different metal mixtures applied within the U0 and U10 condition. In contrast, a reduced growth inhibition is present in the U2 condition after exposure to the metal mixture as compared to the U2-M0 condition.

The potential U-biomarker genes selected after the primer optimization (paragraph 4.1) (primer 33, primer 44, primer 46, primer 48 and primer 52) are further tested in this experiment. The expression of the genes is normalized with the new and existing housekeeping genes. Gene expression levels relative to the control condition (U0-M0) of *L. minor* plants exposed for 7 days to U (0, 2 or 10 μM) and/or metal mixtures (M0, M1, M10 or M100) are given in Figure 5.




Figure 5: Gene expression levels of potential U-biomarker genes for *L. minor* relative to control condition (U0-M0) after exposure for 7 days to U (0, 2 or 10 μM) and/or metal mixtures (M0, M1, M10 or M100). Mean values are given \pm standard error (SE) of at least three biological independent replicates for each condition. Significant differences (two-way ANOVA, p -value < 0.05) within the U0 condition are given with small letters, within the U2 condition with capital letters and within the U10 condition with small letters and an apostrophe. Significant differences (two-way ANOVA, p -value < 0.05) between the U0 condition and U2/U10 condition are given with an asterisk.

In the absence of U, no significant effects are observed between the different metal concentrations applied in primer 33. The same trend is present in the U2 condition. In contrast, the U10 condition shows a metal-dependent response, with a significant difference between M0 and M100. Furthermore primer 33 shows a strong U-dependent response with a 2-fold upregulation after exposure to 2 μ M U (non-significant) and a 5-fold upregulation after 10 μ M U (significant) in the absence of the metal mixture. The M0, M1, M10 and M100 of the U10 condition are significantly higher than their respective control without U.

In contrast to our expectations, a significant increase is observed in the expression of primer 44 after exposure to M100 in the absence of U. No significant effect of the metal mixture is observed when U is applied (2 or 10 μ M). There is a significant increase in the gene expression after exposure to U10 as compared to U0 for M0, M1 and M10 metal mixtures and a significant decrease after exposure to U2 as compared to U0 for M1 and M100.

For primer 46, no significant effects are observed between the different metal mixtures within the U0 and U2 concentration. Within the U10 condition, exposure to the M100 concentration induced a significant increase in the transcript level compared to the other metal concentrations. By comparing the different U concentrations within the same metal mixture applied, a significant increase in gene expression is only present in the U10-M100 condition as compared to U0-M100.

No significant effects are present in the transcript levels of primer 48 for the different metal concentrations within one U concentration. After the application of U, there is a significant decrease in the expression of primer 48 for almost all the metal mixtures applied. However, this downregulation is not concentration-dependent. As such, the expression level at U2 almost drops to zero and increases slightly again at U10.

For the different metal mixtures tested for primer 52, no significant changes are observed within one U concentration, except within the U2 condition. The M100 concentration at 2 μ M U had a higher expression level than the other metal concentrations. After exposure to U, a significant downregulation is obtained whereby the expression levels in M0, M1 and M10 of the U2 condition and M0, M10 and M100 of the U10 condition are significant lower than their corresponding control.

All of the primers were further tested in the time kinetics experiment whereby the gene expression is followed in function of time. This gives the opportunity to find fast-reacting biomarkers.

4.4 Time kinetics experiment

During the time kinetics experiment *L. minor* plants were exposed to three concentrations of the metal mixture (M0, M1, M100) in combination with the three U concentrations (0, 2 and 10 μ M). Samples were taken after 1, 4 and 7 days and the growth of the *L. minor* plants is followed by measuring the total area of the plants and counting the number of fronds (Appendix E). The percentage of growth inhibition (%I_r) is given as the FW of the plants linked to the total area. Table 7 gives the %I_r of the time kinetics experiment.

Table 7: Percentage of growth inhibition (%I_r) calculated from the FW of the plants linked to the total area of *L. minor* plants exposed for 1, 4 or 7 days to U (0, 2 or 10 μ M) and/or metal mixtures (M0, M1, M100). The percentage of growth inhibition of the control condition (U0-M0) is 0%. Mean values are given \pm standard error (SE) of at least four biological independent replicates for each condition. Significant differences (two-way ANOVA, *p*-value <0.05) within the U0 condition are given with small letters, within the U2 condition with capital letters and within the U10 condition with small letters and an apostrophe. Significant differences (two-way ANOVA, *p*-value <0.05) between the U0 condition and U2/U10 condition are given with an asterisk.

| | | 0 μ M U | 2 μ M U | 10 μ M U |
|--------------|-------------|----------------------------------|----------------------------------|-----------------------------------|
| Day 1 | M0 | 0.00% \pm 3.37% ^a | -6.51% \pm 8.27% ^A | -3.85% \pm 4.58% ^{a'} |
| | M1 | -8.00% \pm 4.36% ^a | -15.18% \pm 8.45% ^A | 7.38% \pm 6.66% ^{a'} |
| | M100 | -10.01% \pm 5.45% ^a | -13.43% \pm 8.32% ^A | -5.42% \pm 4.12% ^{a'} |
| Day 4 | M0 | 0.00% \pm 7.68% ^a | 7.27% \pm 2.25% ^A | 23.27% \pm 0.85% ^{a'*} |
| | M1 | -5.73% \pm 0.23% ^a | -2.60% \pm 4.35% ^A | 12.21% \pm 0.93% ^{a'*} |
| | M100 | -3.11% \pm 2.47% ^a | 9.08% \pm 2.60% ^A | 15.90% \pm 6.21% ^{a'*} |
| Day 7 | M0 | 0.00% \pm 1.55% ^a | 0.55% \pm 2.65% ^A | 16.54% \pm 2.78% ^{a'*} |
| | M1 | 5.95% \pm 0.63% ^{ab} | 2.43% \pm 1.91% ^A | 21.32% \pm 1.11% ^{a'*} |
| | M100 | 14.69% \pm 2.22% ^b | 3.02% \pm 3.18% ^{A*} | 23.41% \pm 1.29% ^{a'} |

After 1 day of exposure, the different U concentrations and metal mixtures did not significantly affect the %I_r. At day 4, no significant differences are obtained within one U concentration for the different metal mixtures. But exposure to 10 μ M U leads to a significant growth inhibition compared to their respective controls exposed to 0 μ M U. The same trend for the U10 condition is observed at day 7. No significant differences are obtained in the U2 condition for the different metal mixtures. However, plants grown in the absence of U showed a significant growth reduction after exposure to M100 as compared to U0-M0 after 7 days of exposure.

To find fast-reacting biomarkers, the expression levels of the potential U-biomarker genes was measured and normalized with the new and existing housekeeping genes. Gene expression levels relative to the control condition (U0-M0) of *L. minor* plants exposed for 1, 4 or 7 days to U (0, 2 or 10 μ M) and/or metal mixtures (M0, M1, M100) are given in Figure 6 (primer 33), Figure 7 (primer 44), Figure 8 (primer 46), Figure 9 (primer 48) and Figure 10 (primer 52).




Figure 6: Gene expression levels of potential U-biomarker primer 33 for *L. minor* relative to control condition (U0-M0) after exposure for 1, 4 or 7 days to U (0, 2 or 10 μ M) and/or metal mixtures (M0, M1, M100). Mean values are given \pm standard error (SE) of at least three biological independent replicates for each condition. Significant differences (two-way ANOVA, p -value <0.05) within the U0 condition are given with small letters, within the U2 condition with capital letters and within the U10 condition with small letters and an apostrophe. Significant differences (two-way ANOVA, p -value <0.05) between the U0 condition and U2/U10 condition are given with an asterisk.

At day 1 no significant effects are observed for primer 33 in the U0 and U2 condition between the different metal concentrations applied. Within the U10 condition a metal dependent increase in expression is observed, with a significant increase after exposure to M100 as compared to M0. In general no significant differences are obtained between the different U concentrations for the same metal concentration, with the exception of U10-M0 compared to the non-exposed controls. At day 4, no significant effects are observed within one U concentration. A slight non-significant increase in the gene expression of the U2 and U10 is present compared to the U0 condition, with a significant increase in the U10-M100 condition compared to U0-M100. After 7 days of exposure, no significant effects are observed within one U concentration for the different metal concentrations. Exposing the plants to 2 μ M U leads to an increased expression of the genes (approximately 4 times higher than the control) with a significant increase in the U2-M0 condition. In the U10-M0 condition, the transcript level is 60 times higher. When metals are present with 10 μ M U, the expression rises up to 30 times higher for M1 and 20 times higher for M100. The expression levels in all of the U10 conditions are significant increased compared to their corresponding control conditions.




Figure 7: Gene expression levels of potential U-biomarker primer 44 for *L. minor* relative to control condition (U0-M0) after exposure for 1, 4 or 7 days to U (0, 2 or 10 μ M) and/or metal mixtures (M0, M1, M100). Mean values are given \pm standard error (SE) of at least three biological independent replicates for each condition. Significant differences (two-way ANOVA, p -value <0.05) within the U0 condition are given with small letters, within the U2 condition with capital letters and within the U10 condition with small letters and an apostrophe. Significant differences (two-way ANOVA, p -value <0.05) between the U0 condition and U2/U10 condition are given with an asterisk.

At day 1 no significant differences are observed for primer 44 within one U concentration for the different metal mixtures. A significant upregulation in the gene expression is obtained in all the U10 conditions compared to their respective controls in U0. In contrast no significant differences are present in U2 condition compared to U0. At day 4 significant differences are obtained within the U0 and U2 condition for the different metal mixtures. As such, an increased expression is present after exposure to M100 in both conditions. This increase is not shown in the U10 condition. The transcript level of primer 44 is significantly lower after exposure to U2-M0 compared to U0-M0. In contrast, a 10-fold increase in expression is present in the U10 conditions, with a significant upregulation in U10-M1 compared to their corresponding control. At day 7, no significant differences are observed within one U concentration, except for the M100 of U0 which is significant higher than the other metal concentrations for U0. There is significant decrease in the gene expression after exposure to U2 as compared to U0 for M0 and M100 metal mixtures and a significant increase after exposure to U10 as compared to U0 for M0 and M1.




Figure 8: Gene expression levels of potential U-biomarker primer 46 for *L. minor* relative to control condition (U0-M0) after exposure for 1, 4 or 7 days to U (0, 2 or 10 μ M) and/or metal mixtures (M0, M1, M100). Mean values are given \pm standard error (SE) of at least three biological independent replicates for each condition. Significant differences (two-way ANOVA, p -value <0.05) within the U0 condition are given with small letters, within the U2 condition with capital letters and within the U10 condition with small letters and an apostrophe. Significant differences (two-way ANOVA, p -value <0.05) between the U0 condition and U2/U10 condition are given with an asterisk.

At day 1 and 4 no significant differences are observed for primer 46 within one U concentration. In general no significant differences are obtained between the U concentrations for the same metal concentration, with the exception of U10-M100 of day 4 compared to its control. At day 7, no significant differences are observed within the U0 and U2 condition. For the U10 condition, the transcript level in the M0 metal concentration is significantly higher than in the M100 concentration. Between the different U concentrations, the transcript levels in the M0 and M1 of the U10 conditions are significantly upregulated as compared to their controls in the U0 condition.




Figure 9: Gene expression levels of potential U-biomarker primer 48 for *L. minor* relative to control condition (U0-M0) after exposure for 1, 4 or 7 days to U (0, 2 or 10 μ M) and/or metal mixtures (M0, M1, M100). Mean values are given \pm standard error (SE) of at least three biological independent replicates for each condition. Significant differences (two-way ANOVA, p -value <0.05) within the U0 condition are given with small letters, within the U2 condition with capital letters and within the U10 condition with small letters and an apostrophe. Significant differences (two-way ANOVA, p -value <0.05) between the U0 condition and U2/U10 condition are given with an asterisk.

At day 1, almost no significant differences are obtained for primer 48 for the different metal mixtures within one U concentration, except for the significant upregulation at M100 of the U0 condition. Between the U concentrations there is significant increase in gene expression after exposure to U10 as compared to U0 for the M0 and M1 metal mixtures. This in contrast to the significant decrease in gene expression after exposure to both U2 and U10 as compared to U0 for the M100 metal mixture. At day 4 there are no significant differences observed within one U concentration. All the metal concentrations of the U2 and U10 condition show a significant decrease in gene expression as compared to U0. At day 7, a similar trend as at day 4 is observed with no significant differences within one U concentration, except for the expression level in the U2-M100 condition that is significant lower than in U2-M0. In addition, all the metal concentrations of the U2 and U10 condition show a significant decrease in gene expression as compared to U0.




Figure 10: Gene expression levels of potential U-biomarker primer 52 for *L. minor* relative to control condition (U0-M0) after exposure for 1, 4 or 7 days to U (0, 2 or 10 μ M) and/or metal mixtures (M0, M1, M100). Mean values are given \pm standard error (SE) of at least three biological independent replicates for each condition. Significant differences (two-way ANOVA, p -value <0.05) within the U0 condition are given with small letters, within the U2 condition with capital letters and within the U10 condition with small letters and an apostrophe. Significant differences (two-way ANOVA, p -value <0.05) between the U0 condition and U2/U10 condition are given with an asterisk.

At day 1 no effects are observed for primer 52 within one U concentration and between the different U concentrations. At day 4 there are no significant differences within the U0 and U2 condition for the different metal mixtures. In contrast, a significant upregulation in transcript level is present for the U10-M100 condition compared to U10-M0. There is a significant decrease in the gene expression after exposure to U2 as compared to U0 for the M100 metal mixture and a significant decrease after exposure to U10 as compared to U0 for M0. At day 7, no significant effects are obtained within the U0 and U10 condition for the different metal mixtures. In contrast, there is a significant downregulation in gene expression in the U2-M1 condition compared to U2-M100. Between the U concentrations there is a significant decrease in the gene expression after exposure to U2 as compared to U0 for the M0 and M1 metal mixtures, and for the exposure to U10 as compared to U0 for the M1 metal concentration.

4.5 Summary of the gene expression results

All of the gene expression results given in paragraph 4.3 and 4.4 are analyzed on a number of criteria: exposure dose-response relationship, exposure time-response relationship, sensitivity and chemical specificity.

The exposure dose-response relationship is the up- or downregulation of the primer in function of rising U concentrations. If the gene expression is ascending or descending with the U concentration, this criterion is approved. The exposure time-response relationship is the degree to react in function of time. This information will be obtained from the time-kinetics experiment where the changes in relative gene expression are followed in function of time. The earlier a primer shows an up- or downregulation, the better the time-response relationship. The sensitivity is the responding of the primer to a low concentration of U. A reaction at low U concentrations facilitated the search for an U-biomarker. At last there is the chemical specificity, which is the responding of the primer to the metal mixtures. These are the observed significant differences within one U concentration, i.e. a response to the metal mixture. Since the objective of this thesis is to find biomarkers that react specific to U, primers are approved if no significant effects are observed to the metal mixture [103]. Table 8 gives the overview of the potential U-biomarker primers analyzed with the four biomarker criteria. A check mark indicates that the primer is accepted for this specific criterion, a cross indicates rejection.

Table 8: Overview of potential U-biomarker primers analyzed with the four biomarker criteria. A check mark indicates that the primer is accepted for this specific criterion, a cross indicates rejection.

| | Exposure dose-response relationship | Exposure time-response relationship | Sensitivity | Chemical specificity |
|------------------|--|--|--------------------|-----------------------------|
| Primer 33 | ✓ | ✓ | ✓ | ✓ |
| Primer 44 | × | ✓ | × | × |
| Primer 46 | × | × | × | × |
| Primer 48 | × | ✓ | ✓ | ✓ |
| Primer 52 | × | ✓ | ✓ | ✓ |

Primer 33 matches all four criteria. Primer 44 is only approved by the exposure time-response relationship since it has a fast reaction in gene expression just after one day (Figure 7). Primer 46 did not meet any of the criteria. Primer 48 and primer 52 score exactly the same. Only the exposure dose-response relationship gives negative advice.

4.6 Comet assay

For the first time a comet assay on *L. minor* plants was performed within the BIS group to detect DNA damage. Since it was the aim to test the feasibility of this technique on *L. minor* plants and to optimize some of the parameters, the comet assay was performed on control plants and on plants exposed for four hours to 5 mM MMS, which is an inducer of DNA damage. In total three electrophoresis times were tested: 5, 10 and 15 min. Also two washing steps were tested: two washing steps of 5 min and a washing step of 5 min followed by a washing step of 10 min. An overview of the results is given in Table 9.

Table 9: Overview of different electrophoresis times and washing steps for the performed comet assay

| | | OK? |
|-----------------------------|------------------|-----|
| Electrophoresis time | 5 min | ✓ |
| | 10 min | ✓ |
| | 15 min | ✗ |
| Washing steps | 5 min and 5 min | ✓ |
| | 5 min and 10 min | ✓ |

From the results in Table 9, it can be concluded that an electrophoresis time of 5 and 10 min can be used and that both applied washing steps can be used. An electrophoresis time of 15 min does not give optimal results. As shown in Figure 11 there is some visual DNA damage in the control condition. The comets show tails, which indicates that damaged DNA had been migrated during the electrophoresis, which is not ideal. Based on this results the optimization is continued with 5 and 10 min electrophoresis.




Figure 11: Visual comets of control condition at 15 min of electrophoresis

The visual scoring to determine the different levels of DNA damage is applied on both the control condition and the MMS condition, independently of the electrophoresis time and applied washing steps. Figure 12 gives an example of the results of the comet assay of the control and MMS condition. As shown in the figure, the control condition has mostly comets of class 0 and 1 whereas the MMS condition has comets more spread over the different classes. Table 10 gives a summary of the percentages of the classes from both conditions.




Figure 12: Results of comet assay. Left control condition, right MMS condition both performed with 10 min of electrophoresis and two washing steps of 5 min.

Table 10: Overview of results from the visual scoring of the comet assay. The percentage of each class is given of the control condition and the MMS condition.

| Condition | Class | % of each class |
|-----------|-------|-----------------|
| Control | 0 | 62.73 |
| | 1 | 27.59 |
| | 2 | 9.67 |
| | 3 | 0.00 |
| | 4 | 0.00 |
| MMS | 0 | 2.50 |
| | 1 | 29.58 |
| | 2 | 19.58 |
| | 3 | 16.25 |
| | 4 | 32.08 |

In the absence of MMS, most comets are scored within class 0 and 1 and a few in class 2. No comets are scored in class 3 and 4. In contrast, the MMS condition has comets more spread over the different classes, with >30% in the highest class.

5 DISCUSSION

The aim of this study was divided into two. Firstly, there was the validation of the potential U-biomarker gene (primer 33) and the selection and validation of other potential U-biomarker genes for the freshwater plant *L. minor* when they were exposed to U in combination with an environmentally relevant metal mixture. The environmentally relevant metal mixture was based on concentrations that were measured in the Beaverlodge Lake in Canada [11] and the most important co-contaminants (Cu, Ni, Zn and Pb) were selected and used in this research. The selection of the potential U-biomarker genes was done based on available RNA-sequencing data [7]. Secondly, the feasibility of the comet assay on *L. minor* plants was tested and some of the parameters were optimized using MMS as an inducer of DNA damage. The comet assay can be used as a possible biomarker for detecting DNA damage after exposure to radiation or radionuclides.

5.1 Growth effects induced by exposure to U and the metal mixture

Due to the growth problems of the sterile *L. minor* plants, experiments were not performed according to the OECD guidelines. As such, non-sterile *L. minor* plants were used and plants were grown in a day and night program (14h/10h) at respectively 24/18°C, causing a slight reduction in the growth of the plants. However, the growth of the non-exposed plants (U0-M0) of the 7 days exposure experiment still met the requirements by the OECD (guideline 221) [84], i.e. an average specific growth rate higher than 0.275 d⁻¹ (experimental result of 0.298 d⁻¹). By this, the validity of the experiments is achieved. For the time kinetics experiment an average specific growth rate of 0.272 d⁻¹ is obtained. This indicates that the plants did not meet the requirements by the OECD (guideline 221). Therefore, it is important to repeat the time kinetics experiments to validate the results obtained in this project.

During the 7 days exposure experiment and the time kinetics experiment, the growth of *L. minor* is measured. Following to the OECD guidelines, the total area of the plants and the number of fronds is counted. The FW of the plants can be linked to the total area. Those three endpoints do not give the same results. According to Horemans et al. (2016) [99] the frond number is the least reliable growth parameter considering it not being directly related to the total biomass. Under U stress the frond size will decrease more rapidly than the frond number. As a result the most sensitive endpoint is the total area. Within this thesis the percentage of growth inhibition is calculated from the FW linked to the total area.

In all U concentrations used (0, 2 and 10 µM), plants exposed to M1000 metal concentrations have >50% growth inhibition. This indicates high toxicity at these extremely high metal concentrations, which is a thousandfold higher than environmentally relevant concentrations. As the initial goal of this thesis was to find biomarkers that can be used as early warning systems, i.e. limited effects at the cellular and individual level, this high metal concentration will be excluded from further discussion.

The growth of the *L. minor* plants in the 7 days exposure experiment was inhibited with approximately 20-30% when exposed to 2 and 10 µM U in the absence of metals. While these conditions have a limited growth inhibition at 2 µM U and 15% inhibition at 10 µM U in the time kinetics experiment. As such, there seems to be some differences in growth inhibition in the two experiments, which cannot be explained at the moment. The growth inhibition of the time kinetics experiments is similar to the results obtained by Horemans et al. (2016) [99], at which a 30% growth inhibition was obtained at 10 µM U exposure.

Since the results of Horemans et al. were obtained using the OECD guidelines, a real comparison can only be made when the current experiments are repeated under the same growth conditions.

In accordance with previous experiments [12] a hormesis-like effect is obtained in the U2 condition of the 7 days exposure experiment with all of the metal concentrations. This can be explained by the fact that some of the metals in the metal mixture (Cu, Ni and Zn) are essential for the plant growth. In the absence of these metals, the plants experience toxicity of U. Vanhoudt et al. (2011) [44] proved that U induces a reduction of Cu and Zn concentrations in *A. thaliana* plants. Although this reduced element concentration has not been shown before for *L. minor* plants, it seems that adding those metals to the medium reduces the U toxicity. Therefore, a possible hypothesis is that also in *L. minor* plants, U will influence the nutrient composition of the plants leading to a shortage of essential metals. The effects of the 7 days exposure experiment are not present at day 7 of the time kinetics experiment. Currently there are no explanations for these different results, but studying the uptake of the various elements would help to provide an explanation. In the U10 condition, U shows an adversely effect on the growth of *L. minor* with or without the metal mixtures, in both experiments. No hormesis-like effect is observed, which indicates toxicity.

5.2 Gene expression of potential U-biomarkers

Within this project 10 primers for potential U-biomarker genes were developed and their selectivity was tested on available cDNA of a previous U and metal mixture experiment [12] using real-time qPCR. Based on the results in Figure 3, a selection of five potential U-biomarkers was made and the gene expression levels were measured during the 7 days exposure experiment and the time kinetics experiment. All primers were analyzed on four criteria: 1) exposure dose-response relationship; 2) exposure time-response relationship; 3) sensitivity; and 4) chemical specificity (Table 8).

The function of the gene related to primer 46 is "response to oxidative stress". Although the first screening (Figure 3) showed a dose-related increase in transcript levels and a chemical specificity to U, those results could not be validated in further experiments (Figure 5 and Figure 8). As such, primer 46 did not meet any of the criteria which indicates that this primer can probably not be used as an ideal U-biomarker. In addition, the expression levels in both the 7 days exposure experiment and the time kinetics experiment showed a lower upregulation than results obtained from the cDNA from the previous experiment, indicating the need for the repetition of the experiments with sterile *L. minor* plants according to the OECD guidelines.

Primer 44 is linked to a gene with the function "negative regulation of growth". The transcript levels of this gene were already significantly upregulated after 1 day of exposure to U10, although this was not accompanied yet with a reduced growth of the plants (Table 7). Furthermore, the transcript levels remain upregulated at day 4 and day 7, although non-significant, indicating a fast and long-lasting response to U exposure (exposure time-response relationship). This corresponds to the reduced growth of the plants at day 4 and day 7 after exposure to the highest U concentration, although there is no direct relationship between gene expression and higher levels of biological complexity. In contrast no significant differences were present in U2 condition compared to U0, excluding the dose-response relationship and sensitivity criteria. The expression levels of primer 44 in the 7 days exposure experiment and the time kinetics experiment are also much less upregulated compared to the results obtained from the cDNA from a previous experiment after exposure

to U10. In addition, no significant effects of metal exposure were present in the previous experiment, while in the 7 days exposure experiment and the time kinetics experiment a significant upregulation was observed after exposure to U0-M100, excluding the chemical specificity criteria. The increased expression at U0-M100 was followed by a significant reduction in growth at day 7 in this condition, indicating that an increased expression in this gene precedes a reduction in growth. Thus, primer 44 may not be considered as an ideal U-biomarker according to the analysis in Table 8, but it is definitely a fast-reacting gene.

Primer 48 had only negative advice for the exposure dose-response relationship. The gene related to this primer has as function "response to oxidative stress". It is known that U causes oxidative stress in *L. minor* plants, which was shown by an increased activity of different antioxidative enzymes [21]. In addition, Horemans et al. (2014) [21] has shown that also Cd provokes oxidative stress in *L. minor* plants. However, the antioxidative responses to U and Cd are not identical. For example, the antioxidative enzyme guaiacol peroxidase was strongly induced by Cd whereas it was inhibited by U. Therefore, a possible hypothesis is that due to a different antioxidative response, primer 48 is specific towards U stress. However, more research is necessary with other metals in the environmentally metal mixture, including Cd, to further investigate the chemical specificity of this gene.

The gene related to primer 52 has a function in the response to auxin. As for primer 48, only the exposure dose-response relationship gives negative advice for this primer to become an U-biomarker. It has been hypothesized before that U disrupts the transport and gradient of auxin in the root in *A. thaliana* plants [104]. However, also other metals (e.g. Cd) are known to influence the expression of auxin-related genes. Therefore, it would be interesting to include more metals in the metal mixture to further investigate the chemical specificity of this potential U-biomarker.

Primer 33 is the only primer that matches all four criteria, a primer which is related to a gene encoding for a cytochrome P450 family protein. Similar results are obtained in all the experiments after 7 days of exposure (Figure 3, Figure 5 and Figure 6). The expression levels of the U10 conditions varies from approximately a 25-fold upregulation after 10 μ M U in the previous experiment, an 8-fold upregulation in the 7 days exposure experiment and a 60-fold upregulation in the time kinetics experiment. Although the differences in gene expression levels between the different experiments cannot be explained at this moment, it is clear that this gene reacts specifically and consistent to U-exposure. Since this primer was already selected in previous experiments, primer 33 could be validated as biomarker for U exposure in the current experiments. However, a repetition of the experiments according to the OECD guidelines is advised.

5.3 Comet assay optimization on *L. minor*

The comet assay on *L. minor* was performed for the first time within the BIS group to detect DNA damage. The aim was to test the feasibility of this technique on *L. minor* plants and to optimize some of the parameters. From the first results, it can be concluded that an electrophoresis time of 5 or 10 min can be used to perform the comet assay on *L. minor*, and also both washing steps are approved because similar results are obtained after both tested washing steps. An electrophoresis time of 15 min is considered to be too long. This time was only tested on control plants, but all of the cell nuclei were damaged probably due to the long electrophoresis time. They all had a longer comet tail in comparison with the control plants after 5 or 10 min electrophoresis. The visual scoring based on 5 recognizable classes of comets (Table 6) clearly shows induction of DNA damage in the

MMS treated samples, with almost 68% of the nuclei in class 2, 3 and 4. However, this visual scoring requires some training. If this training is not provided, this technique seems to be quite subjective. A better way to score the comets is with the use of a specialized program such as Comet Assay IV (Score comets). By this, more reliable/consistent results are obtained. The comet assay can definitely be used in further investigations on *L. minor* and in the repetition of the 7 days exposure experiment and time kinetics experiment

CONCLUSION

The aim of this research was to validate the potential U-biomarker gene (primer 33) and to select and validate other potential U-biomarker genes for the freshwater plant *L. minor* when they are exposed to U in combination with an environmentally relevant metal mixture of metals (Cu, Ni, Zn and Pb) based on concentrations that were measured in the Beaverlodge Lake in Canada. Primers for potential U-biomarker genes were developed and tested on available cDNA of a previous U and metal mixture experiment. Five primers with an efficiency >90% and that reacted specifically to U were selected and further tested in two experiments, a 7 days exposure experiment and a time kinetics experiment.

In the 7 days exposure experiment *L. minor* plants were exposed for 7 days to a metal mixture in combination with U. In all U concentrations used, plants exposed to M1000 metal concentrations have >50% growth inhibition. This indicates high toxicity at these extremely high metal concentrations. As the initial goal of this thesis was to find biomarkers that can be used as early warning systems, this high metal concentration was excluded from further discussion.

All five primers were also tested in the time kinetics experiment whereby *L. minor* plants were exposed for 1, 4 or 7 days to three U concentrations (0, 2 and 10 μM) and/or three metal concentrations (M0, M1, M100). Fast-reacting biomarkers could be selected within this experiment.

The primers were analyzed on four criteria: 1) exposure dose-response relationship; 2) exposure time-response relationship; 3) sensitivity; and 4) chemical specificity. The earlier discovered primer 33 matches all four criteria and could be validated as biomarker for U exposure from both experiments. In addition, two new U-biomarker genes were selected within this project, i.e. primer 48 and primer 52 that matches three of the four criteria. Both primers reacted specifically to U already after 4 days of exposure and to the lowest tested U concentration (2 μM). The gene related to primer 48 has as function "response to oxidative stress". U causes oxidative stress in *L. minor* plants by an increased activity of different antioxidative enzymes [21]. But also Cd provokes oxidative stress. However, the antioxidative responses to U and Cd are not identical. More research with other metals in the environmentally metal mixture, including Cd, is necessary to further investigate the chemical specificity of this gene. The gene related to primer 52 has as function in the response to auxin. A hypothesis was made for *A. thaliana* plants, that U disrupts the transport and gradient of auxin in the root of the plants [104]. Also other metals, such as Cd, are known to influence the expression of auxin-related genes. The chemical specificity of this potential U-biomarker under U stress must be further investigated.

A comet assay on *L. minor* was also performed within this project as a possible biomarker for DNA damage after exposure to U. The feasibility of this technique was tested on *L. minor* plants exposed to MMS as a positive control for DNA damage and some of the parameters were optimized. An electrophoresis time of 5 or 10 min was approved. In both the control and the MMS condition cell nuclei were observed. Within the MMS condition, DNA damage was perceived with >30% in the highest class of the visual. In contrast, the control condition has most comets within class 0 and 1 and a few in class 2. However, the used visual scoring based on 5 recognizable classes seems to be quite subjective if no training is provided. A specialized program such as Comet Assay IV (Score comets) could give more reliable/consistent results. But the comet assay can definitely be used in further research on *L. minor*.

Future research should focus on further testing the selected genes under different environmental relevant scenarios, such as the addition of Cd in the metal mixture. This gives possibilities to study the chemical specificity of the genes. It could also be interested to study higher levels of biological complexity. Hereby the reaction of the proteins coded by these genes will be investigated to check if the effects are more than just a change in gene expression. A repetition of the experiments according to the OECD guidelines must be accomplished. This will possibly provide an explanation for the different growth results and could be used as validation of primer 48 and primer 52. However, U-biomarkers based on gene expression are not the only solution. A multiparametric approach is better to predict the outcome and is highly recommended in ecotoxicology. The ideal combination includes 'physiological' biomarkers, biomarkers of general stress and more specific biomarkers [105]. So a combination of the U-biomarker genes and the comet assay can be more applicable in various scenarios.

REFERENCES

- [1] SCK-CEN, "60 jaar ervaring in nucleaire wetenschap en technologie," SCK-CEN, Mol, 2012.
- [2] N. Horemans, *Personal communication*, 2019.
- [3] T. De Meutter, Oxidatieve stress en groei-inhibitie geïnduceerd in eendenkroos (*Lemna minor*) blootgesteld aan cadmium of uranium [eindwerk], Geel: Thomas More, Bachelor in de chemie, afstudeerrichting Milieuzorg, 2013.
- [4] A. Van Gompel, Stressresponsen geïnduceerd in eendenkroos (*Lemna minor*) blootgesteld aan uranium en cadmium in combinatie met verschillende lichtniveaus [eindwerk], Geel: Thomas More, Bachelor in de chemie, afstudeerrichting Biochemie, 2014.
- [5] M. Nuyts, Herstel van *Lemna minor* na blootstelling aan uranium, cadmium of gammastraling [eindwerk], Geel: Thomas More, Bachelor in Biomedische Laboratoriumtechnologie, afstudeerrichting Farmaceutische en biologische laboratoriumtechnologie, 2015.
- [6] A. Van Hoeck, N. Horemans, M. Van Hees, R. Nauts, D. Knapen, H. Vandenhove and R. Blust, " β -Radiation Stress Responses on Growth and Antioxidative Defense System in Plants: A Study with Strontium-90 in *Lemna minor*," *International Journal of Molecular Sciences*, vol. 16, pp. 15309-15327, 2015.
- [7] A. Van Hoeck, N. Horemans, P. Monsieurs, H. Xuan Cao and H. B. R. Vandenhove, "The first draft genome of the aquatic model plant *Lemna minor* opens the route for future stress physiology research and biotechnological applications," *Biotechnical Biofuels*, vol. 8, no. 188, pp. 1-13, 2015.
- [8] A. Van Hoeck, N. Horemans, R. Nauts, M. Van Hees, H. Vandenhove and R. Blust, "*Lemna minor* plants chronically exposed to ionising radiation: RNA-seq analysis indicates a dose rate dependent shift from acclimation to survival strategies," *Plant Science*, vol. 257, pp. 84-95, 2017.
- [9] A. Van Hoeck, N. Horemans, M. Van Hees, R. Nauts, D. Knapen, H. Vandenhove and R. Blust, "Characterizing dose response relationships: Chronic gamma radiation in *Lemna minor* induces oxidative stress and altered ploidy level," *Journal of Environmental Radioactivity*, vol. 150, pp. 195-202, 2015.
- [10] P. Cvjetko, S. Tolić, S. Šikić, B. Balen, M. Tkalec, Ž. Vidaković-Cifrek and M. Pavlica, "Effect of copper on the toxicity and genotoxicity of cadmium in duckweed (*Lemna minor* L.)," *Archives of Industrial Hygiene and Toxicology*, vol. 61, pp. 287-296, 2010.
- [11] S. Lofts, L. Fevrier, N. Horemans, R. Giblin, C. Bruggeman and H. Vandenhove, "Assessment of co-contaminant effects on uranium and thorium speciation in freshwater using geochemical modelling," *Journal of Environmental Radioactivity*, vol. 149, pp. 99-109, 2015.
- [12] E. Knaepen, Toxic effects induced in *Lemna minor* by an environmentally relevant mixture of uranium and metals: selection of potential uranium biomarker genes [eindwerk], Hasselt: Master in de biomedische wetenschappen, 2016.

- [13] K. Asaduzzaman, M. U. Khandaker, N. A. B. Baharudin, Y. B. M. Amin, M. S. Farook, D. A. Bradley and O. Mahmoud, "Heavy metals in human teeth dentine: A bio-indicator of metals exposure and environmental pollution," *Chemosphere*, vol. 176, pp. 221-230, 2017.
- [14] L. Attademo, F. Bernardini, R. Garinella and M. T. Compton, "Environmental pollution and risk of psychotic disorders: A review of the science to date," *Schizophrenia Research*, vol. 181, pp. 55-59, 2017.
- [15] M. Javed, I. Ahmad, N. Usmani and M. Ahmad, "Studies on biomarkers of oxidative stress and associated genotoxicity and histopathology in *Channa punctatus* from heavy metal polluted canal," *Chemosphere*, vol. 151, pp. 210-219, 2016.
- [16] A. Jamers, K. Van der Ven, L. Moens, J. Robbens, G. Potters, Y. Guisez, R. Blust and W. De Coen, "Effect of copper exposure on gene expression profiles in *Chlamydomonas reinhardtii* based on microarray analysis," *Aquatic Toxicology*, vol. 80, pp. 249-260, 2006.
- [17] B. Salbu, L. Skipperud and O. C. Lind, "Sources Contributing to Radionuclides in the Environment: With Focus on Radioactive Particles," in *Radionuclides in the Environment: Influence of chemical speciation and plant uptake on radionuclide migration*, Switzerland, Springer International, 2015, pp. 1-36.
- [18] E. Saenen, N. Horemans, N. Vanhoudt, H. Vandenhove, G. Biermans, M. Van Hees, J. Wannijn, J. Vangronsveld and A. Cuypers, "Effects of pH on uranium uptake and oxidative stress responses induced in *Arabidopsis thaliana*," *Environmental Toxicology and Chemistry*, vol. 32, no. 9, pp. 2125-2133, 2013.
- [19] J. Laurette, C. Larue, I. Llorens, D. Jaillard, P.-H. Jouneau, J. Bourguignon and M. Carrière, "Speciation of uranium in plants upon root accumulation and root-to-shoot translocation: A XAS and TEM study," *Environmental and Experimental Botany*, vol. 77, pp. 87-95, 2012.
- [20] J. Laurette, C. Larue, C. Mariet, F. Brisset, H. Khodja, J. Bourguignon and M. Carrière, "Influence of uranium speciation on its accumulation and translocation in three plant species: Oilseed rape, sunflower and wheat," *Environmental and Experimental Botany*, vol. 77, pp. 96-107, 2012.
- [21] N. Horemans, M. Van Hees, A. Van Hoeck, E. Saenen, T. N. R. De Meutter, R. Blust and H. Vandenhove, "Uranium and cadmium provoke different oxidative stress responses in *Lemna minor* L.," *Plant Biology*, vol. 17, pp. 91-100, 2014.
- [22] F. W. Pontius, "Defining a Guideline for Uranium," *Journal - American Water Works Association*, vol. 92, no. 8, pp. 18-24, 2000.
- [23] S. C. Sheppard, M. I. Sheppard, M.-O. Gallerand and B. Sanipelli, "Derivation of ecotoxicity thresholds for uranium," *Journal of Environmental Radioactivity*, vol. 79, pp. 55-83, 2005.
- [24] A. Bleise, P. Danesi and W. Burkart, "Properties, use and health effects of depleted uranium (DU): a general overview," *Journal of Environmental Radioactivity*, vol. 64, pp. 93-112, 2003.
- [25] L. S. Morton, C. V. Evans and G. O. Estes, "Heavy Metals in the Environment," *Journal of Environmental Quality*, vol. 31, pp. 15-162, 2002.

- [26] E. Weir, "Uranium in drinking water, naturally," *Canadian Medical Association Journal*, vol. 170, no. 6, pp. 951-952, 2004.
- [27] N. Vanhoudt, H. Vandenhove, K. Smeets, T. Remans, M. Van Hees, J. Wannijn, J. Vangronsveld and A. Cuypers, "Effects of uranium and phosphate concentrations on oxidative stress related responses induced in *Arabidopsis thaliana*," *Plant Physiology and Biochemistry*, vol. 46, pp. 987-996, 2008.
- [28] S. Pereira, V. Camilleri, M. Floriani, I. Cavalié, J. Garnier-Laplace and C. Adam-Guillermin, "Genotoxicity of uranium contamination in embryonic zebrafish cells," *Aquatic Toxicology*, vol. 109, pp. 11-16, 2012.
- [29] N. Meravi and S. K. Prajapati, "Biomonitoring the Genotoxicity of Heavy Metals/Metalloids Present in Soil Contaminated by Fly Ash from Coal-Fired Thermal Power Plant Using *Tradescantia pallida*," in *Phytoremediation, Management of Environmental Contaminants Volume 1*, Cham, Springer, 2015, pp. 169-176.
- [30] R. Factori, S. M. Leles, G. C. Novakowski and C. L. S. C. T. S. M. Rocha, "Toxicity and genotoxicity of water and sediment from streams on dotted duckweed (*Landoltia punctata*)," *Brazilian Journal of Biology*, vol. 74, no. 4, pp. 769-778, 2014.
- [31] E. Lacaze, A. Devaux, R. Mons, S. Bony, J. Garric, A. Geffard and O. Geffard, "DNA damage in caged *Gammarus fossarum* amphipods: A tool for freshwater genotoxicity assessment," *Environmental Pollution*, vol. 159, pp. 1682-1691, 2011.
- [32] Lin, Ai-jun, X.-h. Zhang, M.-m. Chen and Q. Cao, "Oxidative stress and DNA damages induced by cadmium accumulation," *Journal of Environmental Sciences*, vol. 19, pp. 596-602, 2007.
- [33] S. Radić, D. Stipaničev, P. Cvjetko, I. L. Mikelić, M. M. Rajčić, S. Širac, B. Pevalek-Kozlina and M. Pavlica, "Ecotoxicological assessment of industrial effluent using duckweed (*Lemna minor* L.) as a test organism," *Ecotoxicology*, vol. 19, pp. 216-222, 2010.
- [34] T. Gichner, Z. Patková, J. Száková and K. Demnerová, "Cadmium induces DNA damage in tobacco roots, but no DNA damage, somatic mutations or homologous recombination in tobacco leaves," *Mutation Research*, vol. 559, pp. 49-57, 2004.
- [35] S. Barillet, A. Buet, C. Adam and A. Devaux, "Does uranium exposure induce genotoxicity in the teleostean *Danio Rerio*? First experimental results," *Radioprotection*, vol. 40, no. S1, pp. S175-S181, 2005.
- [36] S. Barillet, C. Adam, O. Palluel and A. Devaux, "Bioaccumulation, oxidative stress, and neurotoxicity in *Danio rerio* exposed to different isotopic compositions of uranium," *Environmental Toxicology and Chemistry*, vol. 26, no. 3, pp. 497-505, 2007.
- [37] Z. Hon, J. Österreicher and L. Navrátil, "Depleted uranium and its effects on humans," *Sustainability*, vol. 7, pp. 4063-4077, 2015.
- [38] O. Neves and M. J. Matias, "Assessment of groundwaters quality and contamination problems ascribed to an abandoned uranium mine (Cunha Baixa region, Central Portugal)," *Environmental Geology*, vol. 53, pp. 1799-1810, 2008.

- [39] G. Angenard, V. Muczynski, H. Goffigny, C. Duquenne, R. Frydman, R. Habert, G. Livera and V. Rouiller-Fabre, "In vitro effects of Uranium on human fetal germ cells," *Reproductive Toxicology*, vol. 31, pp. 470-476, 2011.
- [40] D. Ribera, F. Labrot, G. Tisnerat and J.-F. Narbonne, "Uranium in the Environment: Occurrence, Transfer, and Biological Effects," *Reviews of Environmental Contaminations and Toxicology*, vol. 146, pp. 53-89, 1996.
- [41] L. D. Kraemer and D. Evans, "Uranium bioaccumulation in a freshwater ecosystem: Impact of feeding ecology," *Aquatic Toxicology*, Vols. 124-125, pp. 163-170, 2012.
- [42] N. Vanhoudt, N. Horemans, G. Biermans, E. Saenen, J. Wannijn, R. Nauts, M. Van Hees and H. Vandenhove, "Uranium affects photosynthetic parameters in *Arabidopsis thaliana*," *Environmental and Experimental Botany*, vol. 97, pp. 22-29, 2014.
- [43] H. Vandenhove, A. Cuypers, M. Van Hees, G. Koppen and J. Wannijn, "Oxidative stress reactions induced in beans (*Phaseolus vulgaris*) following exposure to uranium," *Plant Physiology and Biochemistry*, vol. 44, pp. 795-805, 2006.
- [44] N. Vanhoudt, H. Vandenhove, N. Horemans, D. M. Bello, M. Van Hees, J. Wannijn, R. Carleer, J. Vangronsveld and A. Cuypers, "Uranium induced effects on development and mineral nutrition of *Arabidopsis thaliana*," *Journal of Plant Nutrition*, vol. 34, pp. 1940-1956, 2011.
- [45] N. Vanhoudt, H. Vandenhove, N. Horemans, T. Remans, K. Opdenakker, K. Smeets, D. M. Bello, J. Wannijn, M. Van Hees, J. Vangronsveld and A. Cuypers, "Unraveling uranium induced oxidative stress related responses in *Arabidopsis thaliana* seedlings. Part I: responses in the roots," *Journal of Environmental Radioactivity*, vol. 102, pp. 630-637, 2011.
- [46] N. Vanhoudt, A. Cuypers, N. Horemans, T. Remans, K. Opdenakker, K. Smeets, D. M. Bello, M. Havaux, J. Wannijn, M. Van Hees, J. Vangronsveld and H. Vandenhove, "Unraveling uranium induced oxidative stress related responses in *Arabidopsis thaliana* seedlings. Part II: responses in the leaves and general conclusions," *Journal of Environmental Radioactivity*, vol. 102, pp. 638-645, 2011.
- [47] A. Straczek, L. Duquene, D. Wegrzynek, E. Chinea-Cano, J. Wannijn, J. Navez and H. Vandenhove, "Differences in U root-to-shoot translocation between plant species explained by U distribution in roots," *Journal of Environmental Radioactivity*, vol. 101, pp. 258-266, 2010.
- [48] B. J. Alloway, "Uranium," in *Heavy Metals in Soils: Trace Metals and Metalloids in Soils and their Bioavailability, Third Edition*, Dordrecht, Springer Science & Business Media, 2013, pp. 656-577.
- [49] V. N. Jha, R. M. Tripathi, N. K. Sethy and S. K. Sahoo, "Uptake of uranium by aquatic plants growing in fresh water ecosystem around uranium mill tailings pond at Jaduguda, Indi," *Science of the Total Environment*, vol. 539, pp. 175-184, 2016.
- [50] S. Dieng, K. Kyser and L. Goding, "Tectonic history of the North American shield recorded in uranium deposits in the Beaverlodge area, northern Saskatchewan, Canada," *Precambrian Research*, vol. 224, pp. 316-340, 2013.

- [51] R. Liang, G. Chi, K. Ashton, N. Blamey and M. Fayek, "Fluid compositions and P-T conditions of vein-type uranium mineralization in the Beaverlodge uranium district, northern Saskatchewan, Canada," *Ore Geology Reviews*, vol. 80, pp. 460-183, 2017.
- [52] V. Oros and A. M. Tudoran, "Exotoxicological effects of heavy metals on duckweed plants (*Lemna minor*). III. Tests for growth rate reducing by copper and iron," *Scientific Bulletin Series D: Mining, Mineral Processing, Non-Ferrous Metallurgy, Geology and Environmental Engineering*, vol. 26, no. 2, pp. 7-19, 2012.
- [53] V. Oros and A. Toma, "Ecotoxicological effects of heavy metals on duckweed plants (*Lemna minor*). II. Tests for growth rate reducing by the zinc," *Scientific Bulletin Series D: Mining, Mineral Processing, Non-Ferrous Metallurgy, Geology and Environmental Engineering*, vol. 26, no. 1, pp. 15-21, 2012.
- [54] W. Drost, M. Matzke and T. Backhaus, "Heavy metal toxicity to *Lemna minor*: studies on the time dependence of growth inhibition and the recovery after exposure," *Chemosphere*, vol. 67, pp. 36-43, 2007.
- [55] D. Sijm, M. Rikken, E. Rorije, T. Traas, M. McLachlan and W. Peijnenburg, "Transport, accumulation and transformation processes," in *Risk Assessment of Chemicals: An Introduction, Second Edition*, Dordrecht, Springer Science & Business Media, 2007, pp. 73-158.
- [56] N. Khellaf and M. Zerdaoui, "Growth, photosynthesis and respiratory response to copper in *Lemna minor*: a potential use of duckweed in biomonitoring," *Iranian Journal of Environmental Health Science & Engineering*, vol. 7, no. 2, pp. 299-306, 2010.
- [57] K. Oorts, "Copper," in *Heavy Metals in Soils: Trace Metals and Metalloids in Soils and their Bioavailability, Third Edition*, Dordrecht, Springer Science & Business Media, 2013, pp. 367-394.
- [58] B. J. Alloway, "Heavy Metals and Metalloids as Micronutrients for Plants and Animals," in *Heavy Metals in Soils: Trace Metals and Metalloids in Soils and their Bioavailability, Third Edition*, Dordrecht, Springer Science & Business Media, 2013, pp. 195-210.
- [59] E. Van Genderen, W. Adams, R. Cardwell, P. van Sprang, R. Arnold, R. Santore and P. Rodriguez, "An Evaluation of the Bioavailability and Aquatic Toxicity Attributed to Ambient Copper Concentrations in Surface Waters from Several Parts of the World," *Integrated Environmental Assessment and Management*, vol. 4, no. 4, pp. 416-424, 2008.
- [60] C. Gonnelli and G. Renella, "Chromium and Nickel," in *Heavy Metals in Soils: Trace Metals and Metalloids in Soils and their Bioavailability, Third Edition*, Dordrecht, Springer Science & Business Media, 2013, pp. 313-333.
- [61] P. Madoni, "The acute toxicity of nickel to freshwater ciliates," *Environmental Pollution*, vol. 109, pp. 53-59, 2000.
- [62] J. M. Palma, D. K. Gupta and F. J. Corpas, "Metalloenzymes Involved in the Metabolism of Reactive Oxygen Species and Heavy Metal Stress," in *Heavy Metal Stress in Plants*, Berlin Heidelberg, Springer, 2013, pp. 1-17.

- [63] E. Van Genderen, W. Adams, R. Cardwell, J. Jolosin, R. Santore and P. Rodriguez, "An Evaluation of the Bioavailability and Aquatic Toxicity Attributed to Ambient Zinc Concentrations in Fresh Surface Waters from Several Parts of the World," *Integrated Environmental Assessment and Management*, vol. 5, no. 3, pp. 426-434, 2009.
- [64] J. Mertens and E. Smolders, "Zinc," in *Heavy Metals in Soils: Trace Metals and Metalloids in Soils and their Bioavailability, Third Edition*, Dordrecht, Springer Science & Business Media, 2013, pp. 465-493.
- [65] Z. u. Hassan, S. Ali, M. Rizwan, A. Hussain, Z. Akbar, N. Rasool and F. Abbas, "Role of Zinc in Alleviating Heavy Metal Stress," in *Essential Plant Nutrients: Uptake, Use Efficiency, and Management*, Springer, 2017, pp. 351-366.
- [66] M. J. Haydon and C. S. Cobbett, "A Novel Major Facilitator Superfamily Protein at the Tonoplast Influences Zinc Tolerance and Accumulation in Arabidopsis," *Plant Physiology*, vol. 143, no. 4, pp. 1705-1719, 2007.
- [67] T. Satarpai, J. Shiowatana and A. Siripinyanond, "Paper-based analytical device for sampling, on-site preconcentration and detection of ppb lead in water," *Talanta*, vol. 154, pp. 504-510, 2016.
- [68] P. B. Tchounwou, C. G. Yedjou, A. K. Patlolla and D. J. Sutton, "Heavy Metal Toxicity and the Environment," in *Molecular, Clinical and Environmental Toxicology, Volume 3: Environmental Toxicology*, Basel, Springer Science & Business Media, 2012, pp. 133-164.
- [69] A. A. Ensafi, A. K. Far and S. Meghdadi, "Highly selective optical-sensing film for lead(II) determination in water samples," *Journal of Hazardous Materials*, vol. 172, pp. 1069-1075, 2009.
- [70] International Lead Association, "Lead in Aquatic Environments. Understanding the Science," Bravington House, London.
- [71] R. R. Dietert and M. S. Peipenbrink, "Lead and Immune Function," *Critical Reviews in Toxicology*, vol. 36, pp. 359-385, 2006.
- [72] S. Magrisso, S. Belkin and Y. Erel, "Lead Bioavailability in Soil and Soil Components," *Water, air, and soil pollution*, vol. 202, pp. 315-323, 2009.
- [73] E. Steinnes, "Lead," in *Heavy Metals in Soils: Trace Metals and Metalloids in Soils and their Bioavailability, Third Edition*, Dordrecht, Springer Science & Business Media, 2013, pp. 395-410.
- [74] D. K. Gupta, H. G. Huang and F. J. Corpas, "Lead tolerance in plants: strategies for phytoremediation," *Environmental Science and Pollution Research*, vol. 20, pp. 2150-2161, 2013.
- [75] B. Balen, M. Tkalec, S. Šikić, S. Tolić, P. Cvjetko, M. Pavlica and Ž. Vidaković-Cifrek, "Biochemical responses of Lemna minor experimentally exposed to cadmium and zinc," *Ecotoxicology*, vol. 20, pp. 815-826, 2011.
- [76] J. P. Groten, V. J. Feron and J. Sühnel, "Toxicology of simple and complex mixtures," *TRENDS in Pharmacological Sciences*, vol. 22, no. 6, pp. 316-322, 2011.

- [77] C. Vellinger, M. Parant, P. Rousselle and P. Usseglio-Polatera, "Antagonistic toxicity of arsenate and cadmium in a freshwater amphipod (*Gammarus pulex*)," *Exotoxicology*, vol. 21, pp. 1817-1827, 2012.
- [78] A. L. Charles, S. J. Markick and P. Ralph, "Toxicity of uranium and copper individually, and in combination, to a tropical freshwater macrophyte (*Lemna aquinoctialis*)," *Chemosphere*, vol. 62, pp. 1224-1233, 2006.
- [79] N. Cedergreen, A. M. Christensen, A. Kamper, P. Kudsk, S. K. Mathiassen, J. C. Streibig and H. Sørensen, "A review of independent action compared to concentration addition as reference models for mixtures of compounds with different molecular target sites," *Environmental Toxicology and Chemistry*, vol. 27, no. 7, pp. 1621-1632, 2008.
- [80] J. S. Meyer, K. J. Farley and E. R. Garman, "Metal mixtures modeling evaluation project: 1. background," *Environmental Toxicology and Chemistry*, vol. 34, no. 4, pp. 726-740, 2015.
- [81] V. Oros, A. Toma and A. Tudoran, "Preliminary tests on phytotoxicity of heavy metals Cu, Zn, Cd and Fe on aquatic plants of duckweed (*Lemna minor*)," *Scientific Bulletin Series C: Fascicle Mechanics, Tribology, Machine Manufacturing Technology*, vol. 25, no. 2, pp. 25-32, 2011.
- [82] M. Tkalec, K. Malaric and B. Pevalek-Kozlina, "Influence of 400, 900, and 1900 MHz Electromagnetic Fields on *Lemna minor* Growth and Peroxidase Activity," *Bioelectromagnetics*, vol. 26, pp. 185-193, 2005.
- [83] J. Pratas, P. J. C. Favas, C. Paulo, N. Rodrigues and M. N. V. Prasad, "Uranium accumulation by aquatic plants from uranium-contaminated water in central Portugal," *International Journal of Phytoremediation*, vol. 14, pp. 221-234, 2012.
- [84] Organisatie voor Economische Samenwerking en Ontwikkeling (OECD), "Revised proposal for a new guideline 221: *Lemna* sp. Growth Inhibition Test," in *OECD guidelines for the testing of chemicals*, 2002.
- [85] P. Lindström-Seppä, K. Urban, U. Honkalampi-Hämäläinen and S. Roy, "Biochemical Responses in Aquatic Plants as Markers of Environmental Contamination," *Alternatives to laboratory animals: ATLA*, vol. 29, pp. 277-282, 2001.
- [86] L. R. Shugart, "DNA Damage as a Biomarker of Exposure," *Ecotoxicology*, vol. 9, pp. 329-340, 2000.
- [87] R. A. Brain and N. Cedergreen, "Biomarkers in Aquatic Plants: Selection and Utility," *Reviews of Environmental Contamination and Toxicology*, vol. 198, pp. 49-109, 2008.
- [88] S. M. Adams, J. P. Giesy, L. A. Tremblay and C. T. Eason, "The use of biomarkers in ecological risk assessment: recommendations from the Christchurch conference on Biomarkers in Ecotoxicology," *Biomarkers*, vol. 6, no. 1, pp. 1-6, 2001.
- [89] L. S. McCarty and K. R. Munkittrick, "Environmental Biomarkers in Aquatic Toxicology: Fiction, Fantasy, or Functional?," *Human and Ecological Risk Assessment*, vol. 2, no. 2, pp. 268-274, 1996.

- [90] M. Durante, "Cytogenetic biomarkers for exposure to multiple stressors," in *Multiple Stressors: A Challenge for the Future*, Dordrecht, Springer, 2007, pp. 287-293.
- [91] T. A. Akhtar, M. A. Lampi and B. M. Greenberg, "Identification of six differentially expressed genes in response to copper exposure in the aquatic plant *Lemna gibba* (duckweed)," *Environmental Toxicology and Chemistry*, vol. 24, no. 7, pp. 1705-1715, 2005.
- [92] E. Pernot, J. Hall, S. Baatout, M. A. Benotmane, E. Blanchardon, S. Bouffler, H. E. Saghire, M. Gomolka, A. Guertler, M. Harms-Ringdahl, P. Jeggo, M. Kreuzer, D. Laurier, C. Lindholm, R. Mkacher, R. Quitens, K. Rothkamm, L. Sabatier, S. Tapio, F. de Vathaire and E. Cardis, "Ionizing radiation biomarkers for potential use in epidemiological studies," *Mutation Research*, vol. 751, pp. 258-286, 2012.
- [93] J. Zhang, Y. Zhang, W. Liu, X. Quan, S. Chen, H. Zhao, Y. Jin and W. Zhang, "Evaluation of removal efficiency for acute toxicity and genotoxicity on zebrafish in anoxic-oxic process from selected municipal wastewater treatment plants," *Chemosphere*, vol. 90, pp. 2662-2666, 2013.
- [94] T. Gichner, Z. Patková, J. Száková, I. Žnidar and A. Mukherjee, "DNA damage in potato plants induced by cadmium, ethyl methanesulphonate and γ -rays," *Environmental and Experimental Botany*, vol. 62, pp. 113-119, 2008.
- [95] A. R. Collins, "The Comet Assay for DNA Damage and Repair," *Molecular Biotechnology*, vol. 26, pp. 249-261, 2004.
- [96] B. Pourrut, E. Pinelli, V. Celiz Mendiola, J. Silvestre and F. Douay, "Recommendations for increasing alkaline comet assay reliability in plants," *Mutagenesis*, vol. 30, pp. 37-43, 2015.
- [97] S. Radić, D. Stipaničev, P. Cvjetko, M. M. Rajčić, S. Širac, B. Pevalek-Kozlina and M. Pavlica, "Duckweed *Lemna minor* as a tool for testing toxicity and genotoxicity of surface waters," *Ecotoxicology and Environmental Safety*, vol. 74, pp. 182-187, 2011.
- [98] G. Koppen and L. Verschaeve, "The alkaline comet test on plant cells: A new genotoxicity test for DNA strands breaks in *Vicia faba* root cells," *Mutation Research*, vol. 360, pp. 193-200, 1996.
- [99] N. Horemans, M. Van Hees, E. Saenen, A. Van Hoeck, V. Smolders, R. Blust and H. Vandenhove, "Influence of nutrient medium composition on uranium toxicity and choice of the most sensitive growth related endpoint in *Lemna minor*," *Journal of Environmental Radioactivity*, vol. 151, pp. 427-437, 2016.
- [100] T. Remans, E. Keunen, G. J. Bex, K. Smeets, J. Vangronsveld and A. Cuypers, "Reliable Gene Expression Analysis by Reverse Transcription-Quantitative PCR: Reporting and Minimizing the Uncertainty in Data Accuracy," *The Plant Cell*, vol. 26, pp. 3829-3837, 2014.
- [101] A. Azqueta, S. Meier, C. Priestley, K. B. Gutschow, G. Brunborg, J. Sallatte, F. Soussaline and A. Collins, "The influence of scoring method on variability in results obtained with the comet assay," *Mutagenesis*, vol. 26, no. 3, pp. 393-399, 2011.
- [102] S. Langie, *Protocol Repair Assay For Measuring Base Excision Repair*.

- [103] S. Long, C. Svendsen, R. O. D. Shore, T. Pottinger, Y. Allen, K. Thomas and K. Chipman, "Evaluation of potential use of biomarkers as long-term monitoring tools in assessing ecological quality in terrestrial and aquatic environments of the UK," *English Nature Research Reports*, vol. 593, pp. 1-27, 2004.
- [104] N. B. Serre, C. Alban, J. Bourguignon and S. Ravanel, "Uncovering the physiological and cellular effects of uranium on the root system of *Arabidopsis thaliana*," *Environmental and Experimental Botany*, vol. 157, pp. 121-130, 2019.
- [105] L. Ferrat, C. Pergent-Martini and M. Roméo, "Assessment of the use of biomarkers in aquatic plants for the evaluation of environmental quality: application to seagrasses," *Aquatic Toxicology*, vol. 65, pp. 187-204, 2003.







APPENDIX A: COMPOSITION OF THE CONTAMINATION SOLUTION OF THE 7 DAYS EXPOSURE EXPERIMENT AND THE TIME KINETICS EXPERIMENT









Table 11: Exact composition of the contamination solution used in the 7 days exposure experiment and the time kinetics experiment









| Condition | V M10000 stock solution (ml) | V U stock solution (ml) | V P05-medium (ml) | V MES (mL) |
|-----------|------------------------------|-------------------------|-------------------|------------|
| U0-M0 | 0 | 0 | 100 | 1 |
| U0-M1 | 0.01 | 0 | 99.99 | 1 |
| U0-M10 | 0.1 | 0 | 99.90 | 1 |
| U0-M100 | 1 | 0 | 99.00 | 1 |
| U0-M1000 | 10 | 0 | 90.00 | 1 |
| U2-M0 | 0 | 0.02 | 99.98 | 1 |
| U2-M1 | 0.01 | 0.02 | 99.97 | 1 |
| U2-M10 | 0.1 | 0.02 | 99.88 | 1 |
| U2-M100 | 1 | 0.02 | 98.98 | 1 |
| U2-M1000 | 10 | 0.02 | 89.98 | 1 |
| U10-M0 | 0 | 0.1 | 99.90 | 1 |
| U10-M1 | 0.01 | 0.1 | 99.89 | 1 |
| U10-M10 | 0.1 | 0.1 | 99.80 | 1 |
| U10-M100 | 1 | 0.1 | 98.90 | 1 |
| U10-M1000 | 10 | 0.1 | 89.90 | 1 |

APPENDIX B: SPECIFICITY TESTING OF POTENTIAL HOUSEKEEPING GENES FOR *L. MINOR*

Table 12: Specificity testing of potential housekeeping genes for *L. minor* based on real-time qPCR (amplification plot and melt curve) and primer efficiencies (%) (NA = not applicable). All primers with a primer efficiency >90%, one peak in the melting curve and the concentration dependent expression of the dilution series starting from the highest concentration were further tested within this project.

| Primer number | Primer efficiency (%) | Amplification plot | Melting curve |
|---------------|-----------------------|--|---|
| Primer 53 | 103.07 |  |  |
| Primer 54 | NA |  |  |
| Primer 55 | NA |  |  |



| | | | |
|------------------|--------------|--|---|
| <p>Primer 56</p> | <p>97.96</p> |  |  |
| <p>Primer 57</p> | <p>96.28</p> |  |  |
| <p>Primer 58</p> | <p>NA</p> |  |  |
| <p>Primer 59</p> | <p>92.90</p> |  |  |

| | | | |
|-----------|--------|--|---|
| Primer 60 | 93.93 |  |  |
| Primer 61 | 86.30 |  |  |
| Primer 62 | 100.52 |  |  |
| Primer 63 | 143.82 |  |  |

| | | | |
|------------------|---------------|--|--|
| <p>Primer 64</p> | <p>NA</p> | | |
| <p>Primer 65</p> | <p>101.19</p> | | |
| <p>Primer 66</p> | <p>138.75</p> | | |
| <p>Primer 67</p> | <p>NA</p> | | |







Primer 68


NA



APPENDIX C: SPECIFICITY TESTING OF POTENTIAL U-BIOMARKER GENES FOR *L. MINOR*

Table 13: Specificity testing of potential U-biomarker genes for *L. minor* based on real-time qPCR (amplification plot and melt curve) and primer efficiencies (%) (NA = not applicable). All primers with a primer efficiency >90%, one peak in the melting curve and the concentration dependent expression of the dilution series starting from the highest concentration were further tested within this project.



| Primer number | Primer efficiency (%) | Amplification plot | Melting curve |
|---------------|-----------------------|--|---|
| Primer 41 | 97.23 |  |  |
| Primer 42 | 75.47 |  |  |
| Primer 43 | 90.79 |  |  |



| | | | |
|-----------|--------|--|--|
| Primer 48 | 90.44 | | |
| Primer 49 | NA | | |
| Primer 50 | 118.93 | | |
| Primer 51 | 110.18 | | |

Primer 52

97.98



APPENDIX D: GROWTH OF *L. MINOR* PLANTS IN 7 DAYS EXPOSURE EXPERIMENT

Table 14: Growth measurements (total area and number of fronds) of *L. minor* plants at day 0, 4 and 7 in 7 days exposure experiment

| Condition | Day 0 | | Day 4 | | Day 7 | |
|------------|-------------------------------|------------------|-------------------------------|------------------|-------------------------------|------------------|
| | Total area (mm ²) | Number of fronds | Total area (mm ²) | Number of fronds | Total area (mm ²) | Number of fronds |
| U0-M0-a | 37.990 | 13 | 108.595 | 37 | 283.547 | 76 |
| U0-M0-b | 32.652 | 11 | 110.985 | 32 | 273.573 | 66 |
| U0-M0-c | 39.010 | 12 | 123.751 | 34 | 294.527 | 74 |
| U0-M0-d | 33.694 | 9 | 107.830 | 26 | 246.426 | 59 |
| U0-M0-e | 38.403 | 12 | 140.146 | 35 | 326.685 | 74 |
| U0-M0-f | 35.465 | 12 | 140.065 | 38 | 326.848 | 76 |
| U0-M1-a | 39.580 | 13 | 136.389 | 33 | 342.587 | 78 |
| U0-M1-b | 38.315 | 14 | 123.249 | 39 | 310.609 | 80 |
| U0-M1-c | 36.214 | 12 | 108.544 | 31 | 311.444 | 76 |
| U0-M1-d | 39.203 | 15 | 161.345 | 46 | 383.685 | 88 |
| U0-M10-a | 31.297 | 12 | 120.972 | 31 | 307.545 | 72 |
| U0-M10-b | 30.225 | 9 | 99.111 | 26 | 227.154 | 50 |
| U0-M10-c | 34.417 | 10 | 122.559 | 31 | 282.028 | 65 |
| U0-M10-d | 36.448 | 14 | 128.441 | 36 | 324.491 | 75 |
| U0-M100-a | 37.030 | 13 | 113.383 | 35 | 262.652 | 76 |
| U0-M100-b | 31.614 | 11 | 106.329 | 31 | 248.923 | 68 |
| U0-M100-c | 32.828 | 11 | 113.168 | 29 | 235.286 | 63 |
| U0-M100-d | 34.178 | 10 | 113.204 | 28 | 234.256 | 63 |
| U0-M1000-a | 38.998 | 13 | 55.448 | 22 | 70.973 | 34 |
| U0-M1000-b | 37.019 | 12 | 51.246 | 23 | 63.747 | 33 |
| U0-M1000-c | 31.895 | 10 | 44.805 | 16 | 54.826 | 25 |
| U0-M1000-d | 37.334 | 14 | 49.154 | 20 | 60.286 | 30 |

| | | | | | | |
|------------|--------|----|---------|----|---------|----|
| U2-M0-a | 35.297 | 12 | 89.136 | 31 | 166.802 | 64 |
| U2-M0-b | 29.913 | 10 | 66.646 | 24 | 133.978 | 51 |
| U2-M0-c | 33.726 | 12 | 85.585 | 27 | 180.523 | 57 |
| U2-M0-d | 36.386 | 11 | 90.106 | 34 | 204.485 | 68 |
| U2-M1-a | 34.087 | 11 | 130.638 | 36 | 329.801 | 76 |
| U2-M1-b | 38.194 | 11 | 120.453 | 35 | 290.320 | 73 |
| U2-M1-c | 35.989 | 10 | 106.873 | 32 | 246.996 | 64 |
| U2-M1-d | 40.983 | 12 | 110.645 | 34 | 304.928 | 73 |
| U2-M10-a | 45.572 | 11 | 155.195 | 34 | 385.397 | 78 |
| U2-M10-b | 38.628 | 13 | 129.951 | 40 | 323.422 | 80 |
| U2-M10-c | 35.983 | 9 | 126.551 | 27 | 354.121 | 62 |
| U2-M10-d | 32.463 | 9 | 116.844 | 28 | 307.260 | 62 |
| U2-M100-a | 35.629 | 9 | 108.131 | 27 | 274.349 | 63 |
| U2-M100-b | 30.417 | 12 | 111.576 | 34 | 312.452 | 76 |
| U2-M100-c | 35.274 | 11 | 127.282 | 30 | 328.770 | 69 |
| U2-M100-d | 32.505 | 12 | 116.794 | 32 | 281.766 | 67 |
| U2-M1000-a | 37.218 | 9 | 57.664 | 22 | 96.502 | 43 |
| U2-M1000-b | 33.427 | 10 | 55.212 | 23 | 89.873 | 40 |
| U2-M1000-c | 35.529 | 11 | 59.450 | 26 | 99.219 | 51 |
| U2-M1000-d | 34.511 | 12 | 57.997 | 31 | 81.804 | 45 |
| U10-M0-a | 34.247 | 13 | 94.004 | 28 | 193.699 | 54 |
| U10-M0-b | 29.850 | 9 | 63.853 | 22 | 128.737 | 43 |
| U10-M0-c | 29.478 | 11 | 69.262 | 27 | 126.405 | 46 |
| U10-M0-d | 31.643 | 12 | 74.270 | 28 | 147.527 | 52 |
| U10-M1-a | 33.669 | 13 | 83.013 | 31 | 167.590 | 53 |
| U10-M1-b | 37.728 | 11 | 107.597 | 31 | 225.677 | 60 |
| U10-M1-c | 33.004 | 12 | 92.044 | 29 | 193.413 | 54 |
| U10-M1-d | 34.793 | 11 | 92.138 | 33 | 183.757 | 59 |
| U10-M10-a | 38.013 | 14 | 102.978 | 34 | 209.783 | 59 |

| | | | | | | |
|-------------|--------|----|---------|----|---------|----|
| U10-M10-b | 34.244 | 9 | 84.773 | 25 | 148.753 | 47 |
| U10-M10-c | 36.258 | 11 | 99.114 | 28 | 196.432 | 54 |
| U10-M10-d | 43.689 | 14 | 129.580 | 42 | 252.552 | 66 |
| U10-M100-a | 24.291 | 9 | 54.722 | 18 | 98.767 | 33 |
| U10-M100-b | 34.262 | 9 | 80.938 | 25 | 164.025 | 49 |
| U10-M100-c | 29.672 | 11 | 76.487 | 25 | 144.005 | 43 |
| U10-M100-d | 38.341 | 12 | 111.254 | 29 | 202.471 | 61 |
| U10-M1000-a | 26.992 | 10 | 42.672 | 23 | 58.852 | 34 |
| U10-M1000-b | 35.610 | 13 | 55.269 | 29 | 89.314 | 43 |
| U10-M1000-c | 42.205 | 11 | 69.859 | 27 | 112.423 | 49 |
| U10-M1000-d | 35.000 | 11 | 57.765 | 27 | 83.784 | 40 |

APPENDIX E: GROWTH OF *L. MINOR* PLANTS IN TIME KINETICS EXPERIMENT

Table 15: Growth measurements (total area and number of fronds) of *L. minor* plants (sampled at day 1) at day 0 and 1 in time kinetics experiment

| Condition | Day 0 | | Day 1 | |
|-----------|-------------------------------|------------------|-------------------------------|------------------|
| | Total area (mm ²) | Number of fronds | Total area (mm ²) | Number of fronds |
| U0-M0-a | 38.645 | 12 | 47.436 | 13 |
| U0-M0-b | 42.230 | 14 | 55.664 | 17 |
| U0-M0-c | 30.760 | 9 | 38.576 | 13 |
| U0-M0-d | 34.460 | 10 | 44.288 | 13 |
| U0-M0-e | 36.618 | 10 | 47.181 | 13 |
| U0-M0-f | 28.211 | 8 | 35.199 | 10 |
| U0-M0-g | 34.458 | 10 | 41.602 | 14 |
| U0-M0-h | 35.372 | 10 | 44.656 | 14 |
| U0-M0-i | 41.774 | 12 | 54.062 | 16 |
| U0-M0-j | 44.797 | 12 | 57.662 | 18 |
| U0-M0-k | 31.558 | 9 | 38.446 | 11 |
| U0-M0-l | 37.098 | 11 | 47.840 | 12 |
| U0-M1-a | 45.147 | 14 | 59.223 | 16 |
| U0-M1-b | 34.924 | 10 | 45.730 | 11 |
| U0-M1-c | 35.673 | 11 | 47.038 | 14 |
| U0-M1-d | 49.410 | 15 | 62.184 | 20 |
| U0-M1-e | 42.327 | 12 | 52.873 | 16 |
| U0-M1-f | 44.052 | 13 | 58.067 | 14 |
| U0-M1-g | 34.907 | 10 | 43.862 | 13 |
| U0-M100-a | 45.078 | 14 | 60.228 | 17 |
| U0-M100-b | 38.387 | 12 | 47.915 | 16 |
| U0-M100-c | 39.517 | 13 | 51.850 | 17 |
| U0-M100-d | 41.678 | 12 | 54.735 | 15 |

| | | | | |
|-----------|--------|----|--------|----|
| U0-M100-e | 44.090 | 12 | 54.174 | 18 |
| U0-M100-f | 30.868 | 10 | 40.535 | 12 |
| U0-M100-g | 41.652 | 13 | 54.761 | 14 |
| U2-M0-a | 41.705 | 12 | 56.361 | 15 |
| U2-M0-b | 44.858 | 11 | 59.744 | 15 |
| U2-M0-c | 49.923 | 13 | 60.431 | 17 |
| U2-M0-d | 44.113 | 13 | 58.856 | 16 |
| U2-M0-e | 34.623 | 11 | 45.116 | 12 |
| U2-M0-f | 35.247 | 10 | 42.433 | 12 |
| U2-M0-g | 43.208 | 11 | 54.534 | 15 |
| U2-M1-a | 37.002 | 11 | 47.711 | 14 |
| U2-M1-b | 47.699 | 11 | 65.215 | 20 |
| U2-M1-c | 43.847 | 14 | 57.481 | 16 |
| U2-M1-d | 39.193 | 13 | 52.297 | 14 |
| U2-M1-e | 34.009 | 10 | 46.755 | 14 |
| U2-M1-f | 47.436 | 13 | 56.403 | 16 |
| U2-M1-g | 46.091 | 15 | 60.675 | 19 |
| U2-M100-a | 41.325 | 13 | 57.863 | 15 |
| U2-M100-b | 46.084 | 13 | 59.769 | 18 |
| U2-M100-c | 43.160 | 11 | 56.291 | 14 |
| U2-M100-d | 44.025 | 12 | 58.445 | 15 |
| U2-M100-e | 30.934 | 9 | 37.549 | 12 |
| U2-M100-f | 40.961 | 12 | 55.405 | 15 |
| U2-M100-g | 39.233 | 11 | 49.006 | 14 |
| U10-M0-a | 37.399 | 12 | 47.621 | 15 |
| U10-M0-b | 38.076 | 11 | 50.807 | 15 |
| U10-M0-c | 43.354 | 12 | 55.969 | 17 |
| U10-M0-d | 46.816 | 13 | 60.261 | 16 |
| U10-M0-e | 41.036 | 13 | 52.316 | 16 |
| U10-M0-f | 30.374 | 9 | 37.705 | 12 |

| | | | | |
|------------|--------|----|--------|----|
| U10-M0-g | 47.957 | 13 | 59.124 | 16 |
| U10-M1-a | 37.604 | 10 | 46.925 | 13 |
| U10-M1-b | 36.961 | 13 | 44.480 | 15 |
| U10-M1-c | 37.373 | 10 | 47.690 | 12 |
| U10-M1-d | 30.163 | 10 | 35.648 | 13 |
| U10-M1-e | 43.245 | 13 | 56.284 | 17 |
| U10-M1-f | 43.749 | 13 | 56.447 | 16 |
| U10-M1-g | 35.948 | 14 | 43.299 | 15 |
| U10-M100-a | 39.735 | 12 | 50.216 | 16 |
| U10-M100-b | 32.053 | 9 | 40.64 | 11 |
| U10-M100-c | 36.192 | 10 | 46.783 | 13 |
| U10-M100-d | 34.342 | 11 | 43.081 | 14 |
| U10-M100-e | 40.593 | 10 | 53.628 | 13 |
| U10-M100-f | 41.767 | 10 | 52.105 | 13 |
| U10-M100-g | 44.897 | 13 | 59.268 | 16 |

Table 16: Growth measurements (total area and number of fronds) of *L. minor* plants (sampled at day 4) at day 0 and 4 in time kinetics experiment

| Condition | Day 0 | | Day 4 | |
|-----------|-------------------------------|------------------|-------------------------------|------------------|
| | Total area (mm ²) | Number of fronds | Total area (mm ²) | Number of fronds |
| U0-M0-a | 35.462 | 13 | 110.848 | 34 |
| U0-M0-b | 40.221 | 11 | 119.838 | 32 |
| U0-M0-c | 39.186 | 9 | 107.017 | 25 |
| U0-M0-d | 36.285 | 11 | 131.712 | 30 |
| U0-M0-e | 41.389 | 8 | 83.468 | 21 |
| U0-M0-f | 32.460 | 10 | 102.674 | 26 |
| U0-M1-a | 41.188 | 11 | 108.949 | 32 |
| U0-M1-b | 39.974 | 11 | 123.076 | 29 |
| U0-M1-c | 32.723 | 10 | 101.077 | 25 |
| U0-M1-d | 40.981 | 13 | 125.547 | 34 |
| U0-M100-a | 38.077 | 10 | 117.813 | 27 |
| U0-M100-b | 41.441 | 13 | 130.784 | 32 |
| U0-M100-c | 31.841 | 9 | 92.787 | 24 |
| U0-M100-d | 40.108 | 12 | 113.035 | 28 |
| U2-M0-a | 49.443 | 13 | 130.572 | 34 |
| U2-M0-b | 40.706 | 11 | 116.471 | 23 |
| U2-M0-c | 36.201 | 10 | 96.757 | 26 |
| U2-M0-d | 40.030 | 11 | 102.216 | 26 |
| U2-M1-a | 44.582 | 13 | 135.932 | 33 |
| U2-M1-b | 36.798 | 9 | 98.539 | 23 |
| U2-M1-c | 38.033 | 10 | 126.808 | 28 |
| U2-M1-d | 39.184 | 10 | 112.979 | 25 |
| U2-M100-a | 39.666 | 10 | 99.009 | 24 |
| U2-M100-b | 43.908 | 12 | 112.134 | 33 |
| U2-M100-c | 43.977 | 14 | 124.573 | 34 |

| | | | | |
|------------|--------|----|---------|----|
| U2-M100-d | 39.002 | 10 | 103.167 | 25 |
| U10-M0-a | 35.183 | 11 | 77.643 | 26 |
| U10-M0-b | 37.560 | 12 | 85.684 | 29 |
| U10-M0-c | 30.195 | 9 | 69.489 | 22 |
| U10-M0-d | 36.732 | 11 | 82.818 | 24 |
| U10-M1-a | 39.969 | 12 | 100.567 | 26 |
| U10-M1-b | 32.646 | 9 | 68.531 | 21 |
| U10-M1-c | 38.308 | 11 | 99.344 | 29 |
| U10-M1-d | 31.559 | 9 | 79.507 | 23 |
| U10-M100-a | 43.554 | 11 | 104.572 | 29 |
| U10-M100-b | 39.766 | 11 | 110.332 | 28 |
| U10-M100-c | 32.750 | 9 | 67.103 | 22 |
| U10-M100-d | 40.885 | 12 | 107.054 | 30 |

Table 17: Growth measurements (total area and number of fronds) of *L. minor* plants (sampled at day 7) at day 0, 4 and 7 in time kinetics experiment

| Condition | Day 0 | | Day 4 | | Day 7 | |
|-----------|-------------------------------|------------------|-------------------------------|------------------|-------------------------------|------------------|
| | Total area (mm ²) | Number of fronds | Total area (mm ²) | Number of fronds | Total area (mm ²) | Number of fronds |
| U0-M0-a | 39.566 | 10 | 125.635 | 30 | 336.561 | 69 |
| U0-M0-b | 33.151 | 9 | 111.078 | 25 | 300.449 | 66 |
| U0-M0-c | 38.830 | 10 | 134.340 | 31 | 362.986 | 75 |
| U0-M0-d | 31.610 | 12 | 113.007 | 33 | 324.381 | 82 |
| U0-M0-e | 32.113 | 9 | 107.216 | 24 | 258.433 | 58 |
| U0-M0-f | 38.535 | 11 | 121.310 | 27 | 358.259 | 75 |
| U0-M1-a | 35.868 | 11 | 100.131 | 25 | 279.348 | 63 |
| U0-M1-b | 39.714 | 11 | 114.549 | 29 | 323.787 | 75 |
| U0-M1-c | 30.456 | 11 | 84.528 | 26 | 247.468 | 61 |
| U0-M1-d | 36.700 | 9 | 104.170 | 24 | 283.789 | 62 |
| U0-M100-a | 33.150 | 8 | 100.748 | 25 | 234.099 | 62 |
| U0-M100-b | 31.571 | 10 | 91.670 | 27 | 217.478 | 65 |
| U0-M100-c | 42.407 | 11 | 114.217 | 29 | 240.993 | 73 |
| U0-M100-d | 37.837 | 11 | 106.558 | 28 | 252.673 | 63 |
| U2-M0-a | 36.343 | 11 | 115.509 | 28 | 363.913 | 72 |
| U2-M0-b | 41.672 | 11 | 118.884 | 27 | 338.230 | 63 |
| U2-M0-c | 33.365 | 10 | 104.276 | 26 | 326.692 | 66 |
| U2-M0-d | 35.968 | 10 | 96.876 | 24 | 290.136 | 58 |
| U2-M1-a | 40.674 | 10 | 131.801 | 29 | 395.376 | 70 |
| U2-M1-b | 32.662 | 9 | 95.781 | 23 | 269.224 | 59 |
| U2-M1-c | 38.112 | 11 | 104.510 | 28 | 307.885 | 68 |
| U2-M1-d | 34.135 | 8 | 95.846 | 22 | 286.781 | 53 |
| U2-M100-a | 35.103 | 10 | 107.651 | 26 | 332.524 | 67 |
| U2-M100-b | 41.965 | 14 | 104.273 | 31 | 293.226 | 73 |
| U2-M100-c | 34.143 | 11 | 101.362 | 26 | 318.465 | 61 |

| | | | | | | |
|------------|--------|----|---------|----|---------|----|
| U2-M100-d | 42.096 | 12 | 120.781 | 33 | 352.153 | 79 |
| U10-M0-a | 38.921 | 11 | 109.693 | 28 | 276.513 | 65 |
| U10-M0-b | 43.266 | 11 | 100.790 | 28 | 255.229 | 60 |
| U10-M0-c | 42.467 | 12 | 101.195 | 28 | 232.440 | 60 |
| U10-M0-d | 36.017 | 12 | 97.437 | 31 | 246.129 | 63 |
| U10-M1-a | 29.125 | 9 | 67.677 | 23 | 162.284 | 51 |
| U10-M1-b | 39.804 | 12 | 98.059 | 24 | 232.670 | 52 |
| U10-M1-c | 40.491 | 13 | 104.923 | 31 | 240.265 | 66 |
| U10-M1-d | 37.399 | 10 | 84.016 | 24 | 199.188 | 53 |
| U10-M100-a | 36.378 | 10 | 87.272 | 26 | 211.715 | 53 |
| U10-M100-b | 41.936 | 11 | 94.021 | 25 | 217.070 | 55 |
| U10-M100-c | 33.779 | 10 | 75.070 | 27 | 185.724 | 53 |
| U10-M100-d | 37.701 | 10 | 80.342 | 26 | 194.800 | 56 |

Auteursrechtelijke overeenkomst

Ik/wij verlenen het wereldwijde auteursrecht voor de ingediende eindverhandeling:
Uranium Toxicity to the Freshwater Plant *Lemna minor* in an Environmentally Relevant Metal Mixture: Selection and Validation of Potential Uranium-Biomarker Genes

Richting: **master in de industriële wetenschappen: biochemie**
Jaar: **2019**

in alle mogelijke mediaformaten, - bestaande en in de toekomst te ontwikkelen - , aan de Universiteit Hasselt.

Niet tegenstaand deze toekenning van het auteursrecht aan de Universiteit Hasselt behoud ik als auteur het recht om de eindverhandeling, - in zijn geheel of gedeeltelijk -, vrij te reproduceren, (her)publiceren of distribueren zonder de toelating te moeten verkrijgen van de Universiteit Hasselt.

Ik bevestig dat de eindverhandeling mijn origineel werk is, en dat ik het recht heb om de rechten te verlenen die in deze overeenkomst worden beschreven. Ik verklaar tevens dat de eindverhandeling, naar mijn weten, het auteursrecht van anderen niet overtreedt.

Ik verklaar tevens dat ik voor het materiaal in de eindverhandeling dat beschermd wordt door het auteursrecht, de nodige toelatingen heb verkregen zodat ik deze ook aan de Universiteit Hasselt kan overdragen en dat dit duidelijk in de tekst en inhoud van de eindverhandeling werd genotificeerd.

Universiteit Hasselt zal mij als auteur(s) van de eindverhandeling identificeren en zal geen wijzigingen aanbrengen aan de eindverhandeling, uitgezonderd deze toegelaten door deze overeenkomst.

Voor akkoord,

Van Dyck, Isabelle

Datum: **6/06/2019**